



CONTRACT NO. 95-309
FINAL REPORT
DECEMBER 1998

**Addendum to
Biogenic Hydrocarbon Inventories
for California:**

Generation of Essential Databases

**Measurements of Biogenic
Hydrocarbons in Ambient Air**

CALIFORNIA ENVIRONMENTAL PROTECTION AGENCY



AIR RESOURCES BOARD
Research Division

MEASUREMENTS OF BIOGENIC HYDROCARBONS IN AMBIENT AIR

Addendum to

**BIOGENIC HYDROCARBON INVENTORIES FOR CALIFORNIA:
GENERATION OF ESSENTIAL DATABASES
Final Report**

Contract No. 95-309

STATE OF CALIFORNIA AIR RESOURCES BOARD

Principal Investigator

Arthur M. Winer

Environmental Science and Engineering Program
Department of Environmental Health Sciences
School of Public Health
University of California
Los Angeles, CA 90095

Co-Investigators

John Karlik
Cooperative Extension
University of California, Bakersfield

Janet Arey
Air Pollution Research Center
University of California, Riverside

Contributing Researchers

Yong Jae Chung
Anni Reissell

December 10, 1998

For more information about the ARB's Research Division,
its research and activities, please visit our Web site:

<http://www.arb.ca.gov/rd/rd.htm>

ACKNOWLEDGEMENTS

The contribution of several individuals from the California Air Resources Board and the South Coast Air Quality Management District, particularly Randy Pasek, Bart Croes and Rudy Eden were gratefully appreciated. Their help was essential before the field study when searching for suitable sampling sites, and their encouragement and input throughout the campaign was much appreciated.

Several persons took part in the demanding task of sampling at the sites, as well as preparations before and during the campaign: Cheryl Harry, Eric Shamansky, Roger Atkinson, Bill Harger, Tricia McElroy, Patricia Phousongphuang, Andrew Chew, and Alvaro Alvarado. Their perseverance and endurance was remarkable and imperative to the success of the study. The ingenuity of Bill Long in the designing and fabrication of the sampling equipment, as well as technical support throughout the campaign, was of utmost importance, especially during the early morning hours at a remote sampling site. The contribution of Sara Aschmann in the sample analysis is gratefully acknowledged. The organizational efforts of Traci Lew, Susan Hackbarth and Susan Inong were invaluable throughout the study.

We thank Alan Bescoby and other members of the Bescoby family for allowing us to perform sampling on their property at Mt. Baldy. We also thank Shed and June Behar for kindly allowing us to perform sampling experiments at their property in Ojai.

We gratefully acknowledge support for this research by the California Air Resources Board.

TABLE OF CONTENTS

| | Page |
|---|------|
| Acknowledgements..... | i |
| Table of Contents..... | ii |
| List of Tables..... | iii |
| List of Figures..... | v |
| Disclaimer..... | 1 |
| 1.0 Introduction and Background..... | 2 |
| 2.0 Sampling Sites and Sampling Times..... | 4 |
| 3.0 Experimental Methods..... | 5 |
| 3.1 Sampling..... | 5 |
| 3.2 Analysis..... | 6 |
| 3.3 Intercomparison for Isoprene..... | 7 |
| 4.0 Results..... | 8 |
| 4.1 Isoprene and Its Atmospheric Reaction Products..... | 8 |
| 4.2 Precision of Sampling and Analysis..... | 8 |
| 4.3 Monoterpenes..... | 20 |
| 5.0 Discussion..... | 38 |
| 5.1 Biogenic Hydrocarbon Emissions..... | 38 |
| 5.2 Isoprene and Its Reaction Products..... | 39 |
| 5.2.1 Methacrolein and Methyl vinyl ketone..... | 40 |
| 5.3 Monoterpenes..... | 48 |
| 5.4 Atmospheric Reactions..... | 49 |
| 6.0 Conclusions and Implications..... | 55 |
| 7.0 References..... | 59 |
| Appendix..... | 63 |

LIST OF TABLES

| <u>Table No.</u> | <u>Page</u> |
|---|-------------|
| 1. Intensive sampling periods and sampling sites..... | 4 |
| 2. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Azusa..... | 9 |
| 3. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Pine Mtn. and Mt. Baldy..... | 11 |
| 4. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Banning..... | 13 |
| 5. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Los Angeles North Main Street..... | 14 |
| 6. Isoprene intercomparison sample results..... | 14 |
| 7. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Azusa..... | 21 |
| 8. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Pine Mtn. and Mt. Baldy..... | 24 |
| 9. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Banning..... | 26 |
| 10. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Los Angeles, North Main Street..... | 28 |
| 11. Rate constants k at 298 K and 760 Torr total pressure of air for the reaction of selected biogenic hydrocarbons with the OH radical, the NO_3 radical, and O_3 (data from Atkinson, 1997 unless noted otherwise)..... | 50 |
| 12. Calculated lifetimes for selected biogenic hydrocarbons with respect to photolysis, reaction with the OH radical, reaction with the NO_3 radical, and reaction with O_3 | 51 |
| A-1.1. Detailed data from measurements of isoprene and its atmospheric reaction products in ambient air at Azusa..... | 63 |
| A-1.2. Detailed data from measurements of isoprene and its atmospheric reaction products in ambient air at Pine Mtn. and Mt. Baldy..... | 69 |

LIST OF TABLES (continued)

| <u>Table No.</u> | | <u>Page</u> |
|---|-----|-------------|
| A-1.3. Detailed data from measurements of isoprene and its atmospheric reaction products in ambient air at Banning..... | 73 | |
| A-1.4. Detailed data from measurements of isoprene and its atmospheric reaction products in ambient air at Los Angeles, North Main Street..... | 77 | |
| B-1.1. Detailed data from measurements of α -pinene, camphene, sabinene, and β -pinene in ambient air at Azusa..... | 78 | |
| B-1.2. Detailed data from measurements of α -pinene, camphene, sabinene, and β -pinene in ambient air at Pine Mtn. and Mt. Baldy..... | 84 | |
| B-1.3. Detailed data from measurements of α -pinene, camphene, sabinene, and β -pinene in ambient air at Banning..... | 88 | |
| B-1.4. Detailed data from measurements of α -pinene, camphene, sabinene, and β -pinene in ambient air at Los Angeles, North Main Street..... | 92 | |
| C-1.1. Detailed data from measurements of limonene, 1,8-cineole and camphor in ambient air at Azusa..... | 93 | |
| C-1.2. Detailed data from measurements of limonene, 1,8-cineole and camphor in ambient air at Pine Mtn. and Mt. Baldy..... | 99 | |
| C-1.3. Detailed data from measurements of limonene, 1,8-cineole and camphor in ambient air at Banning..... | 103 | |
| C-1.4. Detailed data from measurements of limonene, 1,8-cineole and camphor in ambient air at Los Angeles, North Main Street..... | 107 | |

LIST OF FIGURES

| <u>Figure No.</u> | | <u>Page</u> |
|-------------------|---|-------------|
| 1. | Structures of biogenic hydrocarbons measured..... | 3 |
| 2. | Isoprene values during Intensive 2, Aug. 4-6, 1997..... | 15 |
| 3. | Isoprene values during Intensive 3, Aug. 22-23, 1997..... | 16 |
| 4. | Isoprene values during Intensive 4, Sept. 4-7, 1997..... | 17 |
| 5. | Isoprene values during Intensive 5, Sept. 28-29, 1997..... | 18 |
| 6. | Isoprene values during Intensive 6, Oct. 3-4, 1997..... | 19 |
| 7. | Biogenic hydrocarbons at Azusa during Intensive 2, Aug. 4-6, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed are shown..... | 29 |
| 8. | Biogenic hydrocarbons at Azusa during Intensive 3, Aug. 22-23, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed are shown..... | 30 |
| 9. | Biogenic hydrocarbons at Azusa during Intensive 4, Sept. 4-7, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed are shown..... | 31 |
| 10. | Biogenic hydrocarbons at Azusa during Intensive 5, Sept. 28-29, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed are shown..... | 32 |
| 11. | Biogenic hydrocarbons at Azusa during Intensive 6, Oct. 3-4, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed are shown..... | 33 |
| 12. | Biogenic hydrocarbons at Los Angeles, North Main St. during Intensive 3, Aug. 22-23, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed are shown..... | 34 |
| 13. | Biogenic hydrocarbons at Banning during Intensive 4, Sept. 5-6, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed are shown..... | 35 |

LIST OF FIGURES (continued)

| <u>Figure No.</u> | | <u>Page</u> |
|-------------------|--|-------------|
| 14. | Biogenic hydrocarbons at Pine Mtn. during Intensive 4, Sept. 4-7, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed are shown..... | 36 |
| 15. | Biogenic hydrocarbons at Mt. Baldy during Intensive 5, Sept. 28-29, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed are shown..... | 37 |
| 16. | Isoprene, methacrolein and methyl vinyl ketone values at Pine Mtn. during Intensive 2, Aug. 4-6, 1997..... | 41 |
| 17. | Isoprene, methacrolein and methyl vinyl ketone values at Pine Mtn. during Intensive 4, Sept. 4-7, 1997..... | 42 |
| 18. | Isoprene, methacrolein and methyl vinyl ketone values at Mt. Baldy during Intensive 5, Sept. 28-29, 1997..... | 43 |
| 19. | Isoprene, methacrolein and methyl vinyl ketone values at Mt. Baldy during Intensive 6, Oct. 3-4, 1997..... | 44 |
| 20. | Isoprene, methacrolein and methyl vinyl ketone values at Azusa during Intensive 4, Sept. 4-7, 1997..... | 46 |
| 21. | Isoprene, methacrolein and methyl vinyl ketone values at Banning during Intensive 4, Sept. 5-6, 1997..... | 47 |
| 22. | Isoprene, α -pinene, cineole and camphor values at Pine Mtn. during Intensive 2, Aug. 4-6, 1997..... | 53 |
| 23. | Isoprene, α -pinene, cineole and camphor values at Pine Mtn. during Intensive 4, Sept. 4-7, 1997..... | 54 |
| 24. | Isoprene, α -pinene, limonene and cineole values at Mt. Baldy during Intensive 5, Sept. 28-29, 1997..... | 56 |
| 25. | Isoprene, α -pinene, limonene and cineole values at Mt. Baldy during Intensive 6, Oct. 3-4, 1997..... | 57 |

This addendum includes the data originally presented in Chapter 6.0 "MEASUREMENTS OF ISOPRENE AND ITS ATMOSPHERIC REACTION PRODUCTS METHACROLEIN AND METHYL VINYL KETONE IN AMBIENT AIR" of the Final Report for Contract No. 95-309. Additional data is presented here on ambient measurements of the monoterpenes: α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor.

DISCLAIMER


The statement and conclusions in this report are those of the University of California and not necessarily those of the California Air Resources Board. The mention of commercial products, their source, or their use in connection with material reported herein is not to be construed as actual or implied endorsement of such products.

MEASUREMENTS OF BIOGENIC HYDROCARBONS IN AMBIENT AIR


1. Introduction and Background

Volatile organic compounds (VOC) of biogenic origin (BHC, biogenic hydrocarbons) were measured in California's South Coast Air Basin (SoCAB) during the Southern California Ozone Study (SCOS97) conducted in the summer 1997. Anthropogenic hydrocarbons are measured on a routine basis in the SoCAB, but few ambient measurements of biogenic hydrocarbons or their atmospheric reaction products have been conducted in the basin. While on a global scale the dominance of BHC over anthropogenic VOC is estimated to be as much as a factor of 10 (WMO, 1995), U.S. emission inventories put the two sources of VOC at comparable strengths (Guenther et al., 1994) and in urban areas such as the SoCAB anthropogenic VOC clearly dominate (Benjamin et al., 1997). Biogenic hydrocarbons are generally highly reactive in the atmosphere (Carter, 1994; Atkinson, 1997; Atkinson and Arey, 1998) and thus can play an important role in tropospheric chemistry (Trainer et al., 1987; Chameides et al., 1988). Because of their rapid atmospheric reactions, however, the ambient concentrations of BHC are generally low and additional measurements of their atmospheric reaction products are necessary to understand the full impact of BHC on photochemical processes such as ambient ozone formation (Montzka et al., 1993; 1995). For this reason, measurements of isoprene, the BHC emitted in greatest quantity by vegetation (Guenther et al., 1995), and its principal reaction products, methacrolein (MACR) and methyl vinyl ketone (MVK), were undertaken during the SCOS97 campaign. Additionally, the following monoterpenes were measured: α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor. Figure 1 shows the structures of these biogenic hydrocarbons.


The SCOS97 campaign took place during June 16 – October 15, 1997, with six intensive sampling periods and a total of 13 days of intensive sampling. Sampling for biogenic VOC was generally conducted simultaneously at three sites chosen as a mid-basin receptor site for anthropogenic VOC (Azusa), a down-wind receptor site (Banning) and a high elevation site impacted by biogenic VOC (Pine Mountain or Mount Baldy). Solid adsorbent tubes with mass flow controllers were utilized for VOC sample collection





Isoprene




Methyl vinyl ketone





Methacrolein

 α -Pinene


Camphene




Sabinene

 β -Pinene

Limonene



1,8-Cineole



Camphor

Figure 1. Structures of biogenic hydrocarbons measured.

and concentration. After sample collection, the adsorbent tubes were cooled and transported to the laboratory where they were analyzed by gas chromatography with mass selective detection (GC-MSD) using a GC-MSD equipped with an Entech thermal desorption/ preconcentrator unit.

Results for ambient diurnal profiles of isoprene and its atmospheric reaction products, methacrolein and methyl vinyl ketone, as well as α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor for the twelve SCOS97 intensive sampling days between August 4 and October 4 are given in this report.

2. Sampling Sites and Sampling Times

The sampling sites in this study were Azusa, Pine Mountain, Mount Baldy, and Banning. During a single intensive period sampling was also performed at a Los Angeles, North Main Street site. The six intensive sampling periods and sampling sites are presented in Table 1.

Table 1. Intensive sampling periods and sampling sites.

| Sampling Period | Sampling Site |
|-------------------|-------------------------------|
| July 14 | Azusa, Banning |
| August 4 - 6 | Azusa, Banning, Pine Mtn. |
| August 22 - 23 | Azusa, Banning, LA North Main |
| September 4 - 7 | Azusa, Banning, Pine Mtn. |
| September 28 - 29 | Azusa, Mt. Baldy |
| October 3 - 4 | Azusa, Banning, Mt. Baldy |

Azusa represents a mid-basin receptor site east of the main source area of downtown Los Angeles, Pine Mtn. is an elevated site (4539 ft) north of Azusa in the San Gabriel Mountains and Mt. Baldy (sampling at 4000 ft) is situated east of Pine Mtn. The Banning site is located approximately 80 miles east of downtown Los Angeles.

The sampling intervals were 3 hours during daytime and 7-9 hours during nighttime. Typically the sampling periods at Azusa, Pine Mtn., and Mt. Baldy were

0300- 0600, 0600-0900, 0900-1200, 1300-1600, 1600-2000, 2000-0300. VOCs were measured at Azusa during the four daytime sampling periods by other investigators participating in SCOS97. At Banning the sampling periods were 0300-0600, 0600-0900, 0900-1200, 1200-1500, 1500-1800, 1800-0300, chosen to coincide with the schedule of the Photochemical Assessment Monitoring Station (PAMS) VOC sampling conducted by the South Coast Air Quality Management District (SCAQMD) at this site. During August 22 - 23 all sites followed the Banning sampling protocol. During September 4 - 7 nighttime sampling was performed more frequently at Azusa and Pine Mtn. with 3 - 4 hour sampling periods (1700-2000, 2000-2400, 2400-0300, 0300-0600).

3. Experimental Methods

3.1 Sampling

Ambient samples were collected on solid adsorbents housed in borosilicate glass tubes (length 130 mm, outer diameter 6 mm, inner diameter 4 mm). For collection of isoprene, methacrolein, methyl vinyl ketone, and the monoterpenes, the tubes were filled with 250 mg of Carbotrap (20/40 mesh, Supelco, Inc.). Prior to packing, tubes were cleaned by sonicating in methanol and annealed at 480°C. The Carbotrap tubes were conditioned by heating overnight at 350 °C with a constant flow of helium through the tubes. After conditioning and while still warm, the adsorbent tubes were capped with brass nuts, caps (Swagelock) and PTFE ferrules (Alltech Associates, Inc.). The tubes were placed in a cleaned glass jar which was sealed with a metal lid lined with Teflon film. A tube containing activated charcoal (MCB Reagents) was kept in the jar during storage of tubes. Blank samples were treated the same way as samples, excluding actual sampling. Two sampling tubes in series were used, with analyses of the second tube serving to verify that no breakthrough occurred. Jars were stored in a refrigerator at 4 °C before transportation to sampling sites. Tubes were freshly desorbed prior to each intensive sampling period.

To prevent ozone from reacting with labile compounds in the adsorbent during sampling, a scrubber housed in a PTFE compression fitting was placed in front of the first sampling tube (Juttner, 1988; Hoffmann, 1994; Helmig, 1997). The scrubber contained eight copper plies coated with manganese dioxide (Dasibi Corp.). Each scrubber was tested by pumping ozone (250 ppbv) through the scrubber and monitoring the mixing ratio with an

ozone monitor (Model 1003-AH, Dasibi Corp.). The scrubbers were tested before and after each intensive sampling period.

At each site the sampling equipment consisted of a diaphragm pump (Model no. 107CAB 050-TFE, Thomas), mass flow controllers (FC-280SAV and FC-280AV, Tylan General) and a mass flow control unit with four channels (RO-28, Tylan General). The tubes were mounted at a height of 1.8 - 2.0 m (at Azusa the sampling equipment was located on the roof of the SCAQMD station). Each pump, mass flow controller, and control unit was checked for flow rate against a bubble flow meter before and after each intensive sampling period.

Carbotrap samples were collected in duplicate with each sampling tube followed with a back tube. Blank samples transported to the sites were treated the same way as samples, excluding actual sampling. Collected samples were transported to the laboratory as soon as possible for spiking with internal standards, and analysis.

3.2 Analysis

Before analysis the samples were spiked with gaseous internal standards. Isoprene-d₈, fully deuterated isoprene, and tricyclene, a saturated tricyclic compound isomeric with the C₁₀H₁₆ monoterpenes, were utilized as internal standards. Known concentrations of isoprene-d₈ and tricyclene were introduced into a Teflon chamber (7900L) from a vacuum rack with an MKS Baratron 1-100 Torr sensor head and a volume-calibrated Pyrex bulb. After ambient sampling each tube was spiked with 100cc of these standards. External calibration samples were prepared by introducing known amounts of isoprene, isoprene-d₈, methacrolein, methyl vinyl ketone, tricyclene, α-pinene, camphene, β-pinene, limonene, 1,8-cineole and camphor into the chamber, sampling known volumes onto adsorbent tubes, and running calibration samples using the identical method as used for the ambient samples. From these calibration samples, response factors for the compounds against the internal standard compounds were determined with isoprene-d₈ serving as the internal standard for isoprene, methacrolein and methyl vinyl ketone, and tricyclene serving as internal standard for α-pinene, camphene, β-pinene, limonene, 1,8-cineole and camphor. An estimated response factor was used for sabinene quantification.

The analyses were performed with a preconcentrator (Entech 7000, Entech Instruments Inc.) connected to a gas chromatograph (Hewlett Packard, 5890) and a mass selective detector (Hewlett Packard, 5971A). The adsorbed compounds were desorbed in a single tube desorber by heating the Carbotrap tube to 325 °C. Concentration prior to injecting the sample into the gas chromatograph was accomplished using a two-step cryofocusing cold trap dehydration method. The analytical column used for separation of compounds was a DB-5 capillary column (length 60 m, diameter 0.32 mm, phase thickness 1 µm, J&W Scientific) and the oven was programmed from 35 °C to 80 °C at the rate of 4 °C/min, then up to 280 °C at 8 °C/min.

The compounds were identified according to their retention times and selected characteristic ions as determined by running pure gaseous standards adsorbed onto sampling tubes. Compound identifications were previously verified by full mass spectra, operating the mass selective detector in the scanning mode (SCAN). The samples were quantified using the selected ion monitoring (SIM) mode monitoring two or more characteristic ions for each compound and using the areas of the ion chromatograms for quantification. The ion 68 m/z was used for quantification of isoprene (confirming ions 67 m/z, 53 m/z), and 76 m/z for isoprene-d₈. The ion 70 m/z was used for quantification of methacrolein (confirming ions 69 m/z, 43 m/z, 41 m/z) and methyl vinyl ketone (confirming ions 69 m/z, 55 m/z). The ion 93 m/z was used for quantification of tricyclene, α-pinene, camphene, sabinene, and β-pinene (confirming ion 136 m/z); ion 68 m/z for limonene (confirming ions 136 m/z, 93 m/z); ion 154 m/z for 1,8-cineole (confirming ion 68 m/z) and 95 m/z for camphor (confirming ion 152 m/z).

3.3 Intercomparison for Isoprene

On August 1, 1997 isoprene sampling was conducted at the University of California, Riverside, Air Pollution Research Center (APRC) between 1430-1815 to allow an intercomparison between the APRC adsorbent tube sampling and canister sampling. Carbotrap tubes were used to collect 5 simultaneous samples for analysis and quantification of isoprene at APRC, as detailed above. Two canister samples (using precleaned canisters supplied by Desert Research Institute and Biospheric Research Corporation) were co-located with the APRC adsorbent samplers. The canisters were

mailed to Desert Research Institute and Biospheric Research Corporation for analysis. The results of the adsorbent tube sampling are provided below.

4. Results

4.1 Isoprene and Its Atmospheric Reaction Products

Tables 2 through 5 give the average mixing ratios of isoprene, methacrolein, and methyl vinyl ketone for the five intensive SCOS97 sampling periods between August and October, 1997. The replicate sample values are given in the Appendix, Tables A-1.1 through A-1.4. The results are reported in ppbv, calculated for 760 Torr and 0°C. Following Table 5 are figures of the isoprene time-concentration profiles for the three (or in one instance two) sampling sites monitored during each intensive sampling period. Azusa was a sampling site during every sampling period and the isoprene values at Azusa are given in the top graph, plotted with a 0.8 ppbv ordinate scale for each intensive (Figures 2 through 6). Pine Mtn. was monitored along with Azusa during Intensives 2 and 4 (Figures 2 and 4), while Mt. Baldy was monitored during Intensives 5 and 6 (Figures 5 and 6). Reflecting the higher ambient isoprene concentrations at these elevated sites, an ordinate scale value of 2.5 ppbv has been utilized for graphing isoprene at these mountain sites. Isoprene was monitored at the down-wind Banning site during Intensives 2, 3, 4, and 6 and is shown as the lower graph with a 0.3 ppbv ordinate scale (Figures 2, 3, 4 and 6). During Intensive 3, Los Angeles at North Main Street was a monitoring site in addition to Azusa and Banning (Figure 3).

4.2 Precision of Sampling and Analysis

Table 6 gives the results for the five replicate isoprene samples taken during the August 1 Intercomparison Sampling. A 6.3% relative standard deviation in the measured isoprene concentration was obtained for these samples.

Table 2. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Azusa.

| AZUSA | Date | Sampling | Isoprene ppbv | MACR ppbv | MVK ppbv |
|-----------------|-------------|------------|------------------|--------------|-------------|
| Aug 4-6, 1997 | 8/4/97 | 0600-0900 | 0.23 | 0.09 | 0.46 |
| | 8/4/97 | 0907-1206 | 0.52 | 0.47 | 0.77 |
| | 8/4/97 | 1306-1603 | 0.46 | 0.51 | 0.92 |
| | 8/4/97 | 1700-2000 | 0.75 | 0.71 | 1.06 |
| | 8/4/97 | 2006-0300 | 0.16 | 0.34 | 0.59 |
| | 8/5/97 | 0307-0600 | 0.24 | 1.01 | 0.58 |
| | 8/5/97 | 0618-0900 | 0.47 | 0.82 | 1.73 |
| | 8/5/97 | 0907-1202 | 0.66 | 0.74 | 1.61 |
| | 8/5/97 | 1300-1600 | 0.61 | 0.70 | 1.49 |
| | 8/5/97 | 1701-2000 | 0.73 | 0.46 | 0.72 |
| | 8/5- 8/6/97 | 2005-0300 | 0.44 | 0.41 | 1.20 |
| | 8/6/97 | 0306-0600 | 0.23 | 0.62 | 0.90 |
| | 8/6/97 | 0616-0900 | 0.37 | 0.60 | 1.24 |
| | 8/6/97 | 0906-1200 | 0.51 | 0.35 | 0.75 |
| | 8/6/97 | 1300-1600 | 0.51 | 0.44 | 0.81 |
| | 8/6/97 | 1700-2000 | 0.40 | 0.15 | 0.23 |
| Aug 22-23, 1997 | 8/22/97 | 0318-0600* | 0.24 | 0.29 | 0.53 |
| | 8/22/97 | 0610-0900* | 0.21 | 0.45 | 0.77 |
| | 8/22/97 | 0905-1157 | 0.41 | 0.33 | 0.69 |
| | 8/22/97 | 1210-1500* | 0.52 | 0.44 | 0.78 |
| | 8/22/97 | 1504-1800 | 0.45 | 0.69 | 1.07 |
| | 8/22-23/97 | 1815-0300 | 0.30 | 3.34 | 1.52 |
| | 8/23/97 | 0304-0600 | 0.25 | 0.55 | 0.98 |
| | 8/23/97 | 0615-0900* | 0.26 | 0.70 | 0.94 |
| | 8/23/97 | 0903-1200 | 0.51 | 0.53 | 0.74 |
| | 8/23/97 | 1208-1500 | 0.31 | 0.49 | 0.82 |
| | 8/23/97 | 1505-1800* | 0.37 | 0.46 | 0.70 |
| Sep 4-7, 1997 | 9/4/97 | 0300-0600 | 0.28 | 0.89 | 1.29 |
| | 9/4/97 | 0606-0900 | 0.22 | 0.55 | 0.56 |
| | 9/4/97 | 0904-1200 | 0.57 | 0.46 | 1.54 |
| | 9/4/97 | 1300-1600 | 0.35 | 0.62 | 1.00 |
| | 9/4/97 | 1700-2000 | 0.61 | 0.40 | 0.50 |
| | 9/4/97 | 2014-2400 | 0.13 | 0.26 | 0.25 |
| | 9/5/97 | 0006-0300 | 0.06 | 0.17 | 0.26 |
| | 9/5/97 | 0305-0600 | 0.06 | 0.16 | 0.25 |
| | 9/5/97 | 0615-0900* | 0.17 | 0.48 | 0.38 |
| | 9/5/97 | 0903-1200 | 0.43 | 0.31 | 0.68 |
| | 9/5/97 | 1300-1600 | 0.31 | 0.39 | 1.01 |
| | 9/5/97 | 1700-2000 | 0.37 | 0.32 | 0.41 |
| | 9/5/97 | 2006-2400 | 0.18 | 0.58 | 0.57 |
| | 9/6/97 | 0008-0300 | 0.11 | 0.22 | 0.28 |
| | 9/6/97 | 0306-0600 | 0.09 | 0.30 | 0.36 |

Table 2. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Azusa (continued).

| AZUSA | Date | Sampling | Isoprene ppbv | MACR ppbv | MVK ppbv |
|-----------------|------------|------------|------------------|--------------|-------------|
| | 9/6/97 | 0615-0900 | 0.15 | 0.33 | 0.27 |
| | 9/6/97 | 0903-1200 | 0.35 | 0.50 | 0.65 |
| | 9/6/97 | 1300-1600 | 0.29 | 0.44 | 0.88 |
| | 9/6/97 | 1700-1959 | 0.29 | 0.32 | 0.50 |
| | 9/6/97 | 2010-2400 | 0.08 | 0.16 | 0.18 |
| | 9/7/97 | 0005-0300 | 0.11 | 0.21 | 0.23 |
| | 9/7/97 | 0306-0600 | 0.11 | 0.30 | 0.52 |
| | 9/7/97 | 0604-0900 | 0.13 | 0.35 | 0.31 |
| Sep 28-29, 1997 | 9/28/97 | 0301-0600 | 0.15 | 0.28 | 2.31 |
| | 9/28/97 | 0609-0900 | 0.23 | 0.31 | 0.53 |
| | 9/28/97 | 0904-1200 | 0.50 | 0.37 | 0.83 |
| | 9/28/97 | 1300-1600 | 0.42 | 0.41 | 0.68 |
| | 9/28/97 | 1700-2000* | 0.34 | 0.25 | 0.31 |
| | 9/28-29/97 | 2015-0300 | 0.07 | 0.19 | 0.23 |
| | 9/29/97 | 0310-0600 | 0.13 | 0.36 | 0.29 |
| | 9/29/97 | 0616-0858 | 0.30 | 0.44 | 0.65 |
| | 9/29/97 | 0900-1200 | 0.37 | 0.30 | 0.46 |
| | 9/29/97 | 1300-1600 | 0.38 | 0.61 | 0.74 |
| | 9/29/97 | 1700-2000 | 0.44 | 0.22 | 0.40 |
| Oct 3-4, 1997 | 10/3/97 | 0301-0600 | 0.06 | 0.13 | 0.26 |
| | 10/3/97 | 0614-0900 | 0.06 | 0.07 | 0.12 |
| | 10/3/97 | 0904-1200 | 0.12 | 0.10 | 0.22 |
| | 10/3/97 | 1300-1600 | 0.21 | 0.19 | 0.38 |
| | 10/3/97 | 1700-2000 | 0.15 | 0.15 | 0.44 |
| | 10/3-4/97 | 2011-0300 | 0.06 | 0.19 | 0.50 |
| | 10/4/97 | 0304-0600 | 0.12 | 0.18 | 0.99 |
| | 10/4/97 | 0612-0900 | 0.07 | 0.08 | 0.26 |
| | 10/4/97 | 0901-1200 | 0.16 | 0.16 | 0.36 |
| | 10/4/97 | 1300-1600* | 0.24 | 0.29 | 0.64 |
| | 10/4/97 | 1700-2000 | 0.14 | 0.11 | 0.22 |

*Single sample value reported. See Appendix for details.

Table 3. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Pine Mtn. and Mt. Baldy.

| PINE MTN. | Date | Sampling | Isoprene ppbv | MACR ppbv | MVK ppbv |
|---------------|--------------|------------|------------------|--------------|-------------|
| Aug 4-6, 1997 | 8/4/97 | 1710-2000 | 2.21 | 0.91 | 1.58 |
| | 08/4-08/5/97 | 2025-0259 | 0.03 | 0.58 | 1.11 |
| | 8/5/97 | 0314-0559 | 0.04 | 0.14 | 0.25 |
| | 8/5/97 | 0617-0858 | 1.96 | 0.09 | 0.16 |
| | 8/5/97 | 0911-1200 | 1.00 | 0.33 | 0.83 |
| | 8/5/97 | 1302-1600* | 0.59 | 0.46 | 0.71 |
| | 8/5/97 | 1700-2000 | 1.09 | 0.44 | 0.72 |
| | 8/5/97 | 2015-0255 | 0.08 | 0.08 | 0.19 |
| | 8/6/97 | 0310-0600 | 0.03 | 0.03 | 0.07 |
| | 8/6/97 | 0615-0900 | 0.80 | 0.07 | 0.14 |
| | 8/6/97 | 0915-1200 | 0.41 | 0.19 | 0.36 |
| | 8/6/97 | 1300-1600 | 0.62 | 0.59 | 1.03 |
| Sep 4-7, 1997 | 9/4/97 | 1702-2000 | 0.41 | 0.52 | 0.72 |
| | 9/4/97 | 2008-2400 | 0.03 | 0.33 | 0.54 |
| | 9/5/97 | 0010-0338 | 0.02 | 0.10 | 0.16 |
| | 9/5/97 | 0349-0600 | 0.02 | 0.09 | 0.18 |
| | 9/5/97 | 0615-0900 | 0.63 | 0.10 | 0.20 |
| | 9/5/97 | 0908-1200 | 0.93 | 0.21 | 0.38 |
| | 9/5/97 | 1300-1600 | 1.71 | 1.07 | 1.38 |
| | 9/5/97 | 1700-2000 | 0.59 | 0.30 | 0.49 |
| | 9/5/97 | 2007-2355 | 0.03 | 0.26 | 0.39 |
| | 9/6/97 | 0000-0255 | 0.02 | 0.10 | 0.17 |
| | 9/6/97 | 0301-0600 | 0.03 | 0.10 | 0.14 |
| | 9/6/97 | 0610-0900 | 0.48 | 0.08 | 0.16 |
| | 9/6/97 | 0905-1200 | 0.60 | 0.17 | 0.17 |
| | 9/6/97 | 1300-1600 | 1.58 | 0.25 | 0.35 |
| | 9/6/97 | 1700-2000 | 0.55 | 0.33 | 0.54 |
| | 9/6/97 | 2007-2355 | 0.05 | 0.18 | 0.26 |
| | 9/7/97 | 0000-0255 | 0.03 | 0.16 | 0.24 |
| | 9/7/97 | 0300-0555 | 0.05 | 0.10 | 0.15 |
| | 9/7/97 | 0600-0900 | 1.88 | 0.10 | 0.21 |

Table 3. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Pinc Mtn. and Mt. Baldy (continued).

| MT. BALDY | Date | Sampling | Isoprene ppbv | MACR ppbv | MVK ppbv |
|-----------------|------------|------------|------------------|--------------|-------------|
| Sep 28-29, 1997 | 9/28/97 | 0904-1200 | 1.90 | 0.23 | 0.40 |
| | 9/28/97 | 1300-1600 | 1.17 | 0.37 | 0.93 |
| | 9/28/97 | 1700-2000 | 2.27 | 0.40 | 0.85 |
| | 9/28-29/97 | 2005-0300 | 0.08 | 0.24 | 0.47 |
| | 9/29/97 | 0307-0600 | 0.06 | 0.06 | 0.17 |
| | 9/29/97 | 0606-0900 | 0.65 | 0.06 | 0.15 |
| | 9/29/97 | 0903-1200 | 1.84 | 0.27 | 0.63 |
| | 9/29/97 | 1300-1600 | 1.07 | 0.29 | 0.61 |
| | 9/29/97 | 1700-2000 | 1.92 | 0.39 | 0.79 |
| Oct 3-4, 1997 | 10/3/97 | 0619-0900 | 0.42 | 0.02 | 0.12 |
| | 10/3/97 | 0903-1200 | 1.38 | 0.13 | 0.28 |
| | 10/3/97 | 1301-1600 | 1.23 | 0.48 | 1.01 |
| | 10/3/97 | 1703-2003* | 1.29 | 0.37 | 0.72 |
| | 10/3-4/97 | 2010-0302 | 0.05 | 0.13 | 0.26 |
| | 10/4/97 | 0310-0601 | 0.04 | 0.06 | 0.11 |
| | 10/4/97 | 0612-0900 | 0.26 | 0.04 | 0.10 |
| | 10/4/97 | 0903-1200 | 1.53 | 0.12 | 0.25 |
| | 10/4/97 | 1300-1600 | 0.88 | 0.32 | 0.75 |
| | 10/4/97 | 1700-2000 | 1.29 | 0.36 | 0.66 |

*Single sample value reported. See Appendix for details.

Table 4. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Banning.

| BANNING | Date | Sampling | Isoprene ppbv | MACR ppbv | MVK ppbv |
|-----------------|-------------|------------|------------------|--------------|-------------|
| Aug 4-6, 1997 | 8/4/97 | 0600-0855 | 0.25 | 0.14 | 0.33 |
| | 8/4/97 | 0902-1157 | 0.06 | 0.09 | 0.19 |
| | 8/4/97 | no sample | - | - | - |
| | 8/4/97 | 1504-1800 | 0.21 | 0.12 | 0.28 |
| | 8/4-5/1997 | 1807-0255 | 0.08 | 0.35 | 0.68 |
| | 8/5/97 | 0303-0550 | 0.02 | 0.16 | 0.26 |
| | 8/5/97 | 0604-0855 | 0.33 | 0.20 | 0.31 |
| | 8/5/97 | 0901-1153 | 0.08 | 0.07 | 0.15 |
| | 8/5/97 | 1159-1452 | 0.10 | 0.04 | 0.11 |
| | 8/5/97 | 1458-1800 | 0.13 | 0.07 | 0.13 |
| | 8/5- 8/6/97 | 1816-0255 | 0.07 | 0.22 | 0.39 |
| | 8/6/97 | 0301-0550 | 0.01 | 0.04 | 0.10 |
| | 8/6/97 | 0605-0855 | 0.02 | 0.02 | 0.04 |
| | 8/6/97 | 0901-1150 | 0.02 | 0.02 | 0.05 |
| | 8/6/97 | 1155-1455 | 0.03 | 0.02 | 0.06 |
| | 8/6/97 | 1500-1800 | 0.18 | 0.05 | 0.16 |
| Aug 22-23, 1997 | 8/22/97 | 0600-0855 | 0.12 | 0.07 | 0.36 |
| | 8/22/97 | 0900-1155 | 0.27 | 0.10 | 0.26 |
| | 8/22/97 | 1200-1455 | 0.20 | 0.09 | 0.23 |
| | 8/22/97 | 1500-1800 | 0.28 | 0.14 | 0.32 |
| | 8/22-23/97 | 1813-0300 | 0.08 | 2.37 | 2.93 |
| | 8/23/97 | 0303-0555 | 0.01 | 0.16 | 0.42 |
| | 8/23/97 | 0603-0900 | 0.14 | 0.12 | 0.27 |
| | 8/23/97 | 0903-1200 | 0.22 | 0.07 | 0.18 |
| | 8/23/97 | 1202-1500 | 0.20 | 0.05 | 0.12 |
| | 8/23/97 | 1502-1800 | 0.22 | 0.07 | 0.25 |
| Sep 5-6, 1997 | 9/5/97 | 0600-0855 | 0.12 | 0.21 | 0.39 |
| | 9/5/97 | 0900-1155 | 0.19 | 0.09 | 0.21 |
| | 9/5/97 | 1158-1458 | 0.17 | 0.03 | 0.11 |
| | 9/5/97 | 1501-1800 | 0.17 | 0.18 | 0.59 |
| | 9/5-6/97 | 1824-0257 | 0.05 | 0.45 | 0.53 |
| | 9/6/97 | 0308-0550* | 0.01 | 0.29 | 0.29 |
| | 9/6/97 | 0610-0855* | 0.10 | 0.33 | 0.44 |
| | 9/6/97 | 0900-1158 | 0.03 | 0.05 | 0.14 |
| | 9/6/97 | 1202-1458 | 0.12 | 0.04 | 0.11 |
| | 9/6/97 | 1502-1800* | 0.21 | 0.06 | 0.18 |

Table 4. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Banning (continued).

| BANNING | Date | Sampling | Isoprene ppbv | MACR ppbv | MVK ppbv |
|---------------|-----------|-----------|------------------|--------------|-------------|
| Oct 3-4, 1997 | 10/3/97 | 0600-0858 | 0.05 | 0.08 | 0.25 |
| | 10/3/97 | 0900-1158 | 0.06 | 0.07 | 0.14 |
| | 10/3/97 | 1200-1458 | 0.03 | 0.03 | 0.08 |
| | 10/3/97 | 1500-1800 | 0.19 | 0.15 | 0.36 |
| | 10/3-4/97 | 1807-0258 | 0.04 | 0.14 | 0.18 |
| | 10/4/97 | 0300-0553 | 0.04 | 0.18 | 0.32 |
| | 10/4/97 | 0601-0858 | 0.10 | 0.16 | 0.29 |
| | 10/4/97 | 0900-1158 | 0.14 | 0.10 | 0.30 |
| | 10/4/97 | 1200-1458 | 0.12 | 0.10 | 0.28 |
| | 10/4/97 | 1500-1800 | 0.11 | 0.16 | 0.46 |

*Single sample value reported. See Appendix for details.

Table 5. Average mixing ratios for isoprene, methacrolein, and methyl vinyl ketone at Los Angeles, North Main Street.

| LOS ANGELES N. MAIN ST. | Date | Sampling | Isoprene ppbv | MACR ppbv | MVK ppbv |
|----------------------------|------------|-----------|------------------|--------------|-------------|
| Aug 22-23, 1997 | 8/22/97 | 0558-0855 | 0.37 | 0.77 | 0.50 |
| | 8/22/97 | 0900-1155 | 0.11 | 1.26 | 0.98 |
| | 8/22/97 | 1202-1457 | 0.21 | 0.49 | 0.67 |
| | 8/22/97 | 1500-1755 | 0.17 | 0.27 | 0.47 |
| | 8/22-23/97 | 1806-0556 | 0.23 | 0.28 | 0.41 |
| | 8/23/97 | 0605-0857 | 0.21 | 0.44 | 0.44 |
| | 8/23/97 | 0900-1156 | 0.12 | 0.33 | 0.73 |
| | 8/23/97 | 1202-1458 | 0.15 | 0.39 | 0.82 |
| | 8/23/97 | 1500-1800 | 0.21 | 0.27 | 0.50 |

Table 6. Isoprene intercomparison sample results.

| Sample no. | mg/m ³ | ppbv |
|-----------------------|-------------------|------|
| 1 | 15.77 | 5.19 |
| 2 | 14.83 | 4.88 |
| 3 | 16.70 | 5.49 |
| 4 | 16.57 | 5.45 |
| 5 | 17.54 | 5.77 |
| Average | 16.3 | 5.4 |
| Standard dev. (1s) | 1.0 | 0.3 |
| Relative st. dev. (%) | 6.3 | 6.3 |




Figure 2. Isoprene values during Intensive 2, Aug. 4-6, 1997.






Figure 3. Isoprene values during Intensive 3, Aug. 22-23, 1997.








Figure 6. Isoprene values during Intensive 6, Oct. 3-4, 1997.

4.3 Monoterpenes

Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor for the five intensive SCOS97 sampling periods are given in Tables 7 through 10. The replicate sample values are given in the Appendix, Tables B-1.1 through B-1.4 and C-1.1 through C-1.4. The results are reported in ppbv, calculated for 760 Torr and 0°C. The isoprene and monoterpene concentrations measured at Azusa for each of the five intensive sampling periods are shown in Figures 7 through 11 with isoprene, α -pinene and limonene shown individually and the remaining biogenic hydrocarbons shown as their sum. The sum of the BHC at Azusa reached almost 5 ppbv during the 0600-0900 sampling period on Sept. 28, 1997. As may be seen from Figure 10, this high BHC concentration was largely due to limonene which reached 3.8 ppbv during this sampling period. During the early morning hours on Aug. 22 (0300-0900), high limonene concentrations were also observed (Figure 8), suggesting a potential anthropogenic source for this BHC at Azusa. In general, during the evening and early morning hours at Azusa, the monoterpene concentrations equalled or exceeded the isoprene concentrations observed.

At the Los Angeles North Main St. site, monitored only during the Aug. 22-23 intensive (Figure 12), the monoterpenes again dominated over isoprene during the nighttime and early morning sampling periods. The BHC concentrations measured at Banning were lower than at the other sampling sites. The data for Intensive 4 (Sept. 5-6) are shown in Figure 13. Unlike at Azusa and LA North Main, limonene never was the most abundant BHC at Banning.

Representative BHC data for the mountain sites are shown in Figures 14 (Pine Mountain) and 15 (Mt. Baldy). At both mountain sites isoprene clearly dominates the BHC profile and consequently during the nighttime the total BHC concentrations are substantially lower than the daytime values.

Table 7. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Azusa.

| AZUSA | Date | Sampling | α -Pinene ppbv | Camphene ppbv | Sabinene ppbv | β -Pinene ppbv | Limonene ppbv | 1,8-Cineole ppbv | Camphor ppbv |
|-----------------|------------|------------|--------------------------|------------------|------------------|-------------------------|------------------|---------------------|-----------------|
| Aug 4-6, 1997 | 8/4/97 | 0600-0900 | 0.320 | 0.011 | 0.345 | 0.045 | 0.188 | 0.043 | 0.022 |
| | 8/4/97 | 0907-1206 | 0.111 | 0.005 | 0.103 | 0.016 | 0.092 | 0.041 | 0.018 |
| | 8/4/97 | 1306-1603 | 0.028 | 0.003 | 0.016 | 0.008 | 0.080 | 0.022 | 0.011 |
| | 8/4/97 | 1700-2000 | 0.045 | 0.004 | 0.028 | 0.015 | 0.095 | 0.033 | 0.007 |
| | 8/4/97 | 2006-0300 | 0.180 | 0.008 | 0.153 | 0.037 | 0.190 | 0.037 | 0.013 |
| | 8/5/97 | 0307-0600 | 0.377 | 0.021 | 0.168 | 0.102 | 0.568 | 0.076 | 0.032 |
| | 8/5/97 | 0618-0900 | 0.221 | 0.014 | 0.090 | 0.062 | 0.278 | 0.100 | 0.058 |
| | 8/5/97 | 0907-1202 | 0.122 | 0.006 | 0.112 | 0.020 | 0.100 | 0.053 | 0.031 |
| | 8/5/97 | 1300-1600 | 0.033 | 0.004 | 0.019 | 0.011 | 0.081 | 0.031 | 0.019 |
| | 8/5/97 | 1701-2000 | 0.047 | 0.004 | 0.028 | 0.016 | 0.074 | 0.026 | 0.017 |
| 8/5- 8/6/97 | 2005-0300 | 0.298 | 0.013 | 0.262 | 0.050 | 0.251 | 0.053 | 0.022 | |
| | 8/6/97 | 0306-0600 | 0.453 | 0.016 | 0.376 | 0.068 | 0.372 | 0.072 | 0.024 |
| | 8/6/97 | 0616-0900 | 0.349 | 0.017 | 0.206 | 0.063 | 0.455 | 0.077 | 0.035 |
| | 8/6/97 | 0906-1200 | 0.081 | 0.007 | 0.037 | 0.016 | 0.128 | 0.047 | 0.027 |
| | 8/6/97 | 1300-1600 | 0.030 | 0.003 | 0.016 | 0.009 | 0.094 | 0.026 | 0.017 |
| | 8/6/97 | 1700-2000 | 0.032 | 0.002 | 0.015 | 0.010 | 0.054 | 0.016 | 0.006 |
| Aug 22-23, 1997 | 8/22/97 | 0318-0600* | 0.167 | 0.140 | 0.018 | 0.007 | 2.297 | 0.044 | 0.038 |
| | 8/22/97 | 0610-0900* | 0.374 | 0.008 | 0.256 | 0.060 | 0.693 | 0.064 | 0.030 |
| | 8/22/97 | 0905-1157 | 0.105 | 0.013 | 0.016 | 0.017 | 0.148 | 0.046 | 0.021 |
| | 8/22/97 | 1210-1500* | 0.039 | 0.007 | 0.002 | 0.003 | 0.121 | 0.024 | 0.012 |
| | 8/22/97 | 1504-1800 | 0.034 | 0.005 | 0.005 | 0.008 | 0.165 | 0.027 | 0.010 |
| | 8/22-23/97 | 1815-0300 | 0.143 | 0.009 | 0.050 | 0.025 | 0.199 | 0.030 | 0.012 |
| | 8/23/97 | 0304-0600* | 0.437 | 0.011 | 0.261 | 0.067 | 0.475 | 0.051 | 0.019 |
| | 8/23/97 | 0615-0900* | 0.388 | 0.013 | 0.140 | 0.043 | 0.465 | 0.041 | 0.024 |
| | 8/23/97 | 0903-1200* | 0.093 | 0.004 | 0.032 | 0.011 | 0.080 | 0.032 | 0.016 |
| | 8/23/97 | 1208-1500 | 0.016 | 0.005 | 0.003 | 0.002 | 0.043 | 0.020 | 0.010 |
| | 8/23/97 | 1505-1800* | 0.004 | 0.019 | 0.001 | ** | 0.030 | 0.011 | 0.005 |

Table 7. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Azusa (continued).

| AZUSA | Date | Sampling | α -Pinene ppbv | Camphene ppbv | Sabinene ppbv | β -Pinene ppbv | Limonene ppbv | 1,8-Cineole ppbv | Camphor ppbv |
|---------------|--------|------------|--------------------------|------------------|------------------|-------------------------|------------------|---------------------|-----------------|
| Sep 4-7, 1997 | 9/4/97 | 0300-0600 | 0.527 | 0.011 | 0.261 | 0.074 | 0.636 | 0.026 | 0.017 |
| | 9/4/97 | 0606-0900 | 0.214 | 0.006 | 0.034 | 0.023 | 0.291 | 0.031 | 0.036 |
| | 9/4/97 | 0904-1200 | 0.091 | 0.003 | 0.022 | 0.011 | 0.122 | 0.030 | 0.028 |
| | 9/4/97 | 1300-1600 | 0.021 | 0.006 | 0.006 | 0.006 | 0.107 | 0.035 | 0.021 |
| | 9/4/97 | 1700-2000 | 0.042 | 0.012 | 0.009 | 0.010 | 0.278 | 0.032 | 0.011 |
| | 9/4/97 | 2014-2400 | 0.129 | 0.004 | 0.035 | 0.011 | 0.153 | 0.015 | 0.009 |
| | 9/5/97 | 0006-0300 | 0.115 | 0.010 | 0.010 | 0.008 | 0.113 | 0.021 | 0.010 |
| | 9/5/97 | 0305-0600 | 0.284 | 0.005 | 0.152 | 0.030 | 0.285 | 0.029 | 0.019 |
| | 9/5/97 | 0615-0900* | 0.392 | 0.010 | 0.108 | 0.039 | 0.498 | 0.034 | 0.023 |
| | 9/5/97 | 0903-1200 | 0.125 | 0.006 | 0.029 | 0.016 | 0.163 | 0.031 | 0.023 |
| | 9/5/97 | 1300-1600 | 0.018 | 0.002 | 0.003 | 0.003 | 0.129 | 0.016 | 0.007 |
| | 9/5/97 | 1700-2000 | 0.036 | 0.003 | 0.003 | 0.006 | 0.059 | 0.010 | 0.004 |
| | 9/5/97 | 2006-2400 | 0.100 | 0.004 | 0.031 | 0.015 | 0.180 | 0.015 | 0.007 |
| | 9/6/97 | 0008-0300 | 0.282 | 0.007 | 0.169 | 0.021 | 0.283 | 0.019 | 0.013 |
| | 9/6/97 | 0306-0600 | 0.251 | 0.007 | 0.031 | 0.030 | 0.253 | 0.019 | 0.016 |
| | 9/6/97 | 0615-0900 | 0.329 | 0.011 | 0.010 | 0.036 | 0.274 | 0.035 | 0.026 |
| | 9/6/97 | 0903-1200 | 0.101 | 0.004 | 0.024 | 0.013 | 0.064 | 0.033 | 0.023 |
| | 9/6/97 | 1300-1600 | 0.019 | 0.004 | 0.002 | 0.003 | 0.104 | 0.013 | 0.007 |
| | 9/6/97 | 1700-1959 | 0.022 | 0.003 | 0.007 | 0.005 | 0.097 | 0.008 | 0.002 |
| | 9/6/97 | 2010-2400 | 0.042 | 0.003 | 0.015 | 0.007 | 0.097 | 0.010 | 0.007 |
| | 9/7/97 | 0005-0300* | 0.287 | 0.009 | 0.036 | 0.015 | 0.274 | 0.020 | 0.015 |
| | 9/7/97 | 0306-0600 | 0.465 | 0.010 | 0.305 | 0.043 | 0.637 | 0.036 | 0.031 |
| | 9/7/97 | 0604-0900 | 0.385 | 0.010 | 0.138 | 0.033 | 0.672 | 0.045 | 0.038 |

Table 7. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Azusa (continued).

| AZUSA | Date | Sampling | α -Pinene ppbv | Camphene ppbv | Sabinene ppbv | β -Pinene ppbv | Limonene ppbv | 1,8-Cineole ppbv | Camphor ppbv |
|-----------------|------------|-----------|--------------------------|------------------|------------------|-------------------------|------------------|---------------------|-----------------|
| Sep 28-29, 1997 | 9/28/97 | 0301-0600 | 0.252 | 0.024 | 0.026 | 0.054 | 0.402 | 0.077 | 0.037 |
| | 9/28/97 | 0609-0900 | 0.274 | 0.010 | 0.230 | 0.073 | 3.798 | 0.063 | 0.035 |
| | 9/28/97 | 0904-1200 | 0.069 | 0.005 | 0.023 | 0.012 | 0.08 | 0.042 | 0.033 |
| | 9/28/97 | 1300-1600 | 0.024 | 0.002 | 0.015 | 0.008 | 0.061 | 0.038 | 0.026 |
| | 9/28/97 | 1700-2000 | 0.016 | 0.004 | 0.007 | 0.007 | 0.056 | 0.027 | 0.016 |
| | 9/28-29/97 | 2015-0300 | 0.097 | 0.006 | 0.061 | 0.017 | 0.123 | 0.028 | 0.014 |
| | 9/29/97 | 0310-0600 | 0.222 | 0.011 | 0.189 | 0.051 | 0.599 | 0.063 | 0.042 |
| | 9/29/97 | 0616-0858 | 0.178 | 0.044 | 0.012 | 0.015 | 0.669 | 0.044 | 0.030 |
| | 9/29/97 | 0900-1200 | 0.096 | 0.004 | 0.086 | 0.020 | 0.115 | 0.058 | 0.030 |
| | 9/29/97 | 1300-1600 | 0.026 | 0.002 | 0.021 | 0.009 | 0.195 | 0.042 | 0.026 |
| | 9/29/97 | 1700-2000 | 0.009 | 0.026 | 0.004 | 0.003 | 0.195 | 0.029 | 0.021 |
| Oct 3-4, 1997 | 10/3/97 | 0301-0600 | 0.041 | 0.006 | 0.016 | 0.007 | 0.105 | 0.022 | 0.012 |
| | 10/3/97 | 0614-0900 | 0.058 | 0.005 | 0.042 | 0.013 | 0.153 | 0.056 | 0.034 |
| | 10/3/97 | 0904-1200 | 0.033 | 0.005 | 0.012 | 0.007 | 0.098 | 0.038 | 0.041 |
| | 10/3/97 | 1300-1600 | 0.022 | 0.005 | 0.006 | 0.006 | 0.091 | 0.042 | 0.031 |
| | 10/3/97 | 1700-2000 | 0.016 | 0.004 | 0.010 | 0.010 | 0.105 | 0.025 | 0.013 |
| | 10/3-4/97 | 2011-0300 | 0.093 | 0.004 | 0.045 | 0.010 | 0.067 | 0.021 | 0.012 |
| | 10/4/97 | 0304-0600 | 0.023 | 0.131 | 0.006 | 0.006 | 0.122 | 0.025 | 0.041 |
| | 10/4/97 | 0612-0900 | 0.080 | 0.007 | 0.021 | 0.018 | 0.137 | 0.061 | 0.052 |
| | 10/4/97 | 0901-1200 | 0.011 | 0.053 | 0.006 | 0.006 | 0.125 | 0.039 | 0.051 |
| | 10/4/97 | 1300-1600 | 0.014 | 0.010 | 0.010 | 0.006 | 0.049 | 0.021 | 0.018 |
| | 10/4/97 | 1700-2000 | 0.014 | 0.004 | 0.007 | 0.006 | 0.087 | 0.016 | 0.012 |

*Single sample value reported. See Appendix for details.

**Not detected.

Table 8. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Pine Mtn. and Mt. Baldy.

| PINE MTN. | Date | Sampling | α -Pinene ppbv | Camphene ppbv | Sabinene ppbv | β -Pinene ppbv | Limonene ppbv | 1,8-Cineole ppbv | Camphor ppbv |
|---------------|--------------|------------|--------------------------|------------------|------------------|-------------------------|------------------|---------------------|-----------------|
| Aug 4-6, 1997 | 8/4/97 | 1710-2000 | 0.021 | 0.007 | 0.011 | 0.005 | 0.011 | 0.024 | 0.025 |
| | 08/4-08/5/97 | 2025-0259 | 0.001 | 0.005 | 0.001 | 0.0003 | 0.005 | 0.015 | 0.021 |
| | 8/5/97 | 0314-0559 | 0.001 | 0.001 | 0.001 | 0.0003 | 0.002 | 0.006 | 0.008 |
| | 8/5/97 | 0617-0858 | 0.016 | 0.002 | 0.003 | 0.005 | 0.011 | 0.005 | 0.004 |
| | 8/5/97 | 0911-1200 | 0.053 | 0.008 | 0.031 | 0.007 | 0.012 | 0.031 | 0.030 |
| | 8/5/97 | 1302-1600* | 0.038 | 0.011 | 0.009 | 0.007 | 0.012 | 0.047 | 0.055 |
| | 8/5/97 | 1700-2000 | 0.028 | 0.008 | 0.016 | 0.006 | 0.008 | 0.033 | 0.048 |
| | 8/5/97 | 2015-0255 | 0.003 | 0.001 | 0.001 | 0.001 | 0.003 | 0.016 | 0.029 |
| | 8/6/97 | 0310-0600 | 0.005 | 0.001 | 0.001 | 0.002 | 0.004 | 0.004 | 0.006 |
| | 8/6/97 | 0615-0900 | 0.016 | 0.006 | 0.002 | 0.004 | 0.008 | 0.007 | 0.010 |
| | 8/6/97 | 0915-1200 | 0.031 | 0.005 | 0.018 | 0.004 | 0.009 | 0.023 | 0.028 |
| | 8/6/97 | 1300-1600 | 0.040 | 0.011 | 0.015 | 0.007 | 0.011 | 0.056 | 0.058 |
| Sep 4-7, 1997 | | | | | | | | | |
| | 9/4/97 | 1702-2000 | 0.014 | 0.007 | 0.014 | 0.003 | 0.012 | 0.009 | 0.011 |
| | 9/4/97 | 2008-2400 | 0.001 | 0.001 | 0.001 | 0.001 | 0.003 | 0.008 | 0.008 |
| | 9/5/97 | 0010-0338 | 0.001 | 0.008 | 0.001 | 0.002 | 0.004 | ** | 0.003 |
| | 9/5/97 | 0349-0600 | 0.001 | 0.005 | 0.002 | 0.001 | 0.006 | 0.006 | 0.004 |
| | 9/5/97 | 0615-0900 | 0.005 | 0.014 | 0.002 | 0.002 | 0.005 | ** | 0.005 |
| | 9/5/97 | 0908-1200 | 0.029 | 0.003 | 0.013 | 0.002 | 0.008 | 0.009 | 0.013 |
| | 9/5/97 | 1300-1600 | 0.029 | 0.005 | 0.005 | 0.003 | 0.008 | 0.012 | 0.015 |
| | 9/5/97 | 1700-2000 | 0.028 | 0.004 | 0.013 | 0.004 | 0.009 | 0.009 | 0.010 |
| | 9/5/97 | 2007-2355 | 0.001 | ** | 0.001 | ** | ** | 0.006 | 0.007 |
| | 9/6/97 | 0000-0255 | 0.001 | ** | 0.001 | 0.001 | 0.003 | 0.005 | 0.005 |
| | 9/6/97 | 0301-0600 | 0.001 | 0.001 | 0.001 | 0.001 | 0.002 | ** | 0.004 |
| | 9/6/97 | 0610-0900 | 0.004 | 0.002 | 0.001 | 0.001 | 0.003 | 0.006 | 0.006 |
| | 9/6/97 | 0905-1200 | 0.015 | 0.004 | 0.004 | 0.002 | 0.006 | 0.009 | 0.011 |
| | 9/6/97 | 1300-1600 | 0.015 | 0.004 | 0.005 | 0.002 | 0.005 | 0.011 | 0.011 |
| | 9/6/97 | 1700-2000 | 0.018 | 0.005 | 0.005 | 0.003 | 0.006 | 0.009 | 0.010 |
| | 9/6/97 | 2007-2355 | 0.001 | 0.001 | 0.001 | 0.001 | 0.003 | 0.004 | 0.005 |
| | 9/7/97 | 0000-0255 | 0.001 | 0.009 | 0.002 | 0.002 | 0.003 | ** | 0.004 |
| | 9/7/97 | 0300-0555 | 0.001 | 0.001 | 0.002 | 0.002 | 0.003 | 0.009 | 0.003 |
| | 9/7/97 | 0600-0900 | 0.006 | 0.002 | 0.003 | 0.002 | 0.004 | ** | 0.001 |

Table 8. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Pine Mtn. and Mt. Baldy (continued).

| MT. BALDY | Date | Sampling | α -Pinene ppbv | Camphene ppbv | Sabinene ppbv | β -Pinene ppbv | Limonene ppbv | 1,8-Cineole ppbv | Camphor ppbv |
|-----------------|------------|------------|--------------------------|------------------|------------------|-------------------------|------------------|---------------------|-----------------|
| Sep 28-29, 1997 | 9/28/97 | 0904-1200 | 0.208 | 0.008 | 0.023 | 0.060 | 0.056 | 0.014 | 0.013 |
| | 9/28/97 | 1300-1600 | 0.101 | 0.005 | 0.016 | 0.024 | 0.021 | 0.022 | 0.016 |
| | 9/28/97 | 1700-2000 | 0.132 | 0.014 | 0.007 | 0.040 | 0.030 | 0.022 | 0.012 |
| 9/28-29/97 | 2005-0300 | 0.052 | 0.006 | 0.005 | 0.035 | 0.011 | 0.019 | 0.010 | |
| 9/29/97 | 0307-0600 | 0.058 | 0.005 | 0.006 | 0.038 | 0.012 | 0.014 | 0.008 | |
| | 0606-0900 | 0.076 | 0.006 | 0.009 | 0.043 | 0.017 | 0.016 | 0.007 | |
| 9/29/97 | 0903-1200 | 0.083 | 0.012 | 0.004 | 0.015 | 0.030 | 0.020 | 0.011 | |
| 9/29/97 | 1300-1600 | 0.053 | 0.007 | 0.009 | 0.010 | 0.016 | 0.012 | 0.008 | |
| 9/29/97 | 1700-2000 | 0.084 | 0.010 | 0.007 | 0.031 | 0.022 | 0.019 | 0.009 | |
| Oct 3-4, 1997 | | | | | | | | | |
| | 10/3/97 | 0619-0900 | 0.040 | 0.006 | 0.009 | 0.022 | 0.023 | 0.018 | 0.026 |
| | 10/3/97 | 0903-1200 | 0.050 | 0.008 | 0.007 | 0.017 | 0.034 | 0.031 | 0.041 |
| | 10/3/97 | 1301-1600* | 0.050 | 0.007 | 0.011 | 0.010 | 0.010 | 0.023 | 0.019 |
| | 10/3/97 | 1703-2003 | 0.035 | 0.008 | 0.008 | 0.014 | 0.019 | 0.043 | 0.025 |
| 10/3-4/97 | 2010-0302 | 0.020 | 0.003 | 0.003 | 0.015 | 0.005 | 0.012 | 0.011 | |
| 10/4/97 | 0310-0601 | 0.025 | 0.005 | 0.003 | 0.020 | 0.007 | 0.013 | 0.007 | |
| 10/4/97 | 0612-0900* | 0.046 | 0.005 | 0.008 | 0.033 | 0.019 | 0.016 | 0.007 | |
| 10/4/97 | 0903-1200 | 0.063 | 0.009 | 0.003 | 0.011 | 0.020 | 0.015 | 0.008 | |
| 10/4/97 | 1300-1600 | 0.046 | 0.006 | 0.010 | 0.011 | 0.027 | 0.021 | | |
| 10/4/97 | 1700-2000 | 0.039 | 0.007 | 0.006 | 0.013 | 0.018 | 0.035 | 0.024 | |

*Single sample value reported. See Appendix for details.

**Not detected.

Table 9. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Banning.

| BANNING | Date | Sampling | α -Pinene ppbv | Camphene ppbv | Sabinene ppbv | β -Pinene ppbv | Limonene ppbv | 1,8-Cineole ppbv | Camphor ppbv |
|-----------------|-------------|-----------|--------------------------|------------------|------------------|-------------------------|------------------|---------------------|-----------------|
| Aug 4-6, 1997 | 8/4/97 | 0600-0855 | 0.035 | 0.003 | 0.010 | 0.006 | 0.011 | 0.020 | 0.015 |
| | 8/4/97 | 0902-1157 | 0.014 | 0.001 | 0.007 | 0.004 | 0.004 | 0.005 | 0.004 |
| | 8/4/97 | no sample | - | - | - | - | - | - | - |
| | 8/4/97 | 1504-1800 | 0.011 | 0.002 | 0.004 | 0.003 | 0.007 | 0.015 | 0.018 |
| | 8/4-5/1997 | 1807-0255 | 0.008 | 0.001 | 0.002 | 0.003 | 0.006 | 0.022 | 0.009 |
| | 8/5/97 | 0303-0550 | 0.021 | 0.002 | 0.002 | 0.005 | 0.027 | 0.018 | 0.010 |
| | 8/5/97 | 0604-0855 | 0.060 | 0.005 | 0.007 | 0.012 | 0.023 | 0.020 | 0.013 |
| | 8/5/97 | 0901-1153 | 0.032 | 0.001 | 0.002 | 0.010 | 0.005 | 0.005 | 0.007 |
| | 8/5/97 | 1159-1452 | 0.031 | 0.002 | 0.006 | 0.005 | 0.006 | 0.003 | 0.003 |
| | 8/5/97 | 1458-1800 | 0.019 | 0.001 | 0.002 | 0.006 | 0.006 | 0.005 | 0.004 |
| | 8/5- 8/6/97 | 1816-0255 | 0.008 | 0.002 | 0.001 | 0.002 | 0.007 | 0.035 | 0.025 |
| | 8/6/97 | 0301-0550 | 0.004 | 0.001 | 0.003 | 0.002 | 0.003 | 0.010 | 0.010 |
| | 8/6/97 | 0605-0855 | 0.033 | 0.001 | 0.004 | 0.012 | 0.004 | 0.003 | 0.002 |
| | 8/6/97 | 0901-1150 | 0.044 | 0.002 | 0.003 | 0.012 | 0.005 | 0.003 | 0.002 |
| | 8/6/97 | 1155-1455 | 0.047 | 0.001 | 0.005 | 0.017 | 0.005 | 0.003 | 0.002 |
| | 8/6/97 | 1500-1800 | 0.024 | 0.002 | 0.003 | 0.007 | 0.004 | 0.012 | 0.012 |
| Aug 22-23, 1997 | | | | | | | | | |
| | 8/22/97 | 0600-0855 | 0.039 | 0.003 | 0.004 | 0.005 | 0.027 | 0.012 | 0.007 |
| | 8/22/97 | 0900-1155 | 0.019 | 0.002 | 0.002 | 0.002 | 0.008 | 0.012 | 0.007 |
| | 8/22/97 | 1200-1455 | 0.009 | 0.002 | 0.001 | 0.001 | 0.003 | 0.024 | 0.031 |
| | 8/22/97 | 1500-1800 | 0.010 | 0.002 | 0.001 | 0.002 | 0.004 | 0.014 | 0.013 |
| | 8/22-23/97 | 1813-0300 | 0.005 | 0.002 | 0.001 | 0.001 | 0.008 | 0.030 | 0.019 |
| | 8/23/97 | 0303-0555 | 0.001 | 0.002 | 0.002 | -** | 0.003 | 0.017 | 0.015 |
| | 8/23/97 | 0603-0900 | 0.014 | 0.002 | 0.002 | 0.002 | 0.003 | 0.012 | 0.011 |
| | 8/23/97 | 0903-1200 | 0.007 | 0.001 | 0.001 | 0.002 | 0.006 | 0.006 | 0.006 |
| | 8/23/97 | 1202-1500 | 0.004 | 0.001 | 0.0005 | 0.002 | 0.007 | 0.005 | 0.005 |
| | 8/23/97 | 1502-1800 | 0.005 | 0.001 | 0.0005 | 0.001 | 0.002 | 0.009 | 0.012 |

Table 9. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Banning (continued).

| BANNING | Date | Sampling | α -Pinene ppbv | Camphene ppbv | Sabinene ppbv | β -Pinene ppbv | Limonene ppbv | 1,8-Cineole ppbv | Camphor ppbv |
|---------------|------------|------------|--------------------------|------------------|------------------|-------------------------|------------------|---------------------|-----------------|
| Sep 5-6, 1997 | 9/5/97 | 0600-0855 | 0.064 | 0.005 | 0.004 | 0.007 | 0.016 | 0.038 | 0.013 |
| | 9/5/97 | 0900-1155 | 0.010 | 0.002 | 0.001 | 0.001 | 0.003 | 0.013 | 0.005 |
| | 9/5/97 | 1158-1458 | 0.009 | 0.001 | 0.001 | 0.001 | 0.003 | 0.007 | 0.006 |
| | 9/5/97 | 1501-1800 | 0.007 | 0.001 | 0.001 | 0.001 | 0.003 | 0.009 | 0.008 |
| 9/5-6/97 | 1824-0257* | 0.010 | 0.001 | 0.001 | 0.002 | 0.008 | 0.018 | 0.013 | |
| 9/6/97 | 0308-0550* | 0.017 | 0.002 | 0.002 | 0.003 | 0.010 | 0.023 | 0.010 | |
| 9/6/97 | 0610-0855* | 0.012 | 0.010 | 0.002 | 0.002 | 0.021 | 0.033 | 0.007 | |
| 9/6/97 | 0900-1158 | 0.016 | 0.003 | 0.001 | 0.005 | 0.014 | -** | 0.002 | |
| 9/6/97 | 1202-1458 | 0.007 | 0.001 | 0.001 | 0.003 | 0.009 | 0.008 | | |
| 9/6/97 | 1502-1800* | 0.007 | 0.001 | 0.001 | 0.004 | 0.011 | 0.010 | | |
| Oct 3-4, 1997 | | | | | | | | | |
| | 10/3/97 | 0600-0858* | 0.014 | 0.023 | 0.001 | 0.002 | 0.028 | 0.024 | 0.023 |
| | 10/3/97 | 0900-1158 | 0.003 | 0.073 | -** | -** | 0.021 | 0.010 | 0.013 |
| | 10/3/97 | 1200-1458 | 0.003 | 0.049 | 0.001 | -** | 0.016 | -** | 0.004 |
| | 10/3/97 | 1500-1800* | 0.007 | 0.004 | 0.001 | 0.002 | 0.007 | 0.016 | 0.008 |
| 10/3-4/97 | 1807-0258* | 0.009 | 0.010 | 0.001 | 0.002 | 0.032 | 0.032 | 0.024 | |
| 10/4/97 | 0300-0553 | 0.192 | 0.009 | 0.004 | 0.024 | 0.058 | 0.077 | 0.033 | |
| 10/4/97 | 0601-0858 | 0.336 | 0.012 | 0.006 | 0.035 | 0.044 | 0.057 | 0.018 | |
| 10/4/97 | 0900-1158 | 0.018 | 0.003 | 0.001 | 0.003 | 0.009 | 0.026 | 0.013 | |
| 10/4/97 | 1200-1458 | 0.007 | 0.002 | 0.001 | 0.005 | 0.017 | 0.011 | | |
| 10/4/97 | 1500-1800 | 0.006 | 0.003 | 0.001 | 0.001 | 0.016 | 0.013 | 0.008 | |

*Single sample value reported. See Appendix for details.

**Not detected.

Table 10. Average mixing ratios for α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor at Los Angeles, North Main Street.

| LOS ANGELES N. MAIN ST. | Date | Sampling | α -Pinene ppbv | Camphene ppbv | Sabinene ppbv | β -Pinene ppbv | Limonene ppbv | 1,8-Cineole ppbv | Camphor ppbv |
|----------------------------|------------|-----------|--------------------------|------------------|------------------|-------------------------|------------------|---------------------|-----------------|
| Aug 22-23, 1997 | 8/22/97 | 0558-0855 | 0.292 | 0.022 | 0.050 | 0.032 | 0.338 | 0.120 | 0.047 |
| | 8/22/97 | 0900-1155 | 0.036 | 0.027 | 0.004 | 0.003 | 0.120 | 0.053 | 0.045 |
| | 8/22/97 | 1202-1457 | 0.005 | 0.008 | 0.002 | 0.002 | 0.036 | 0.018 | 0.014 |
| | 8/22/97 | 1500-1755 | 0.004 | 0.001 | 0.001 | 0.002 | 0.025 | 0.007 | 0.005 |
| | 8/22-23/97 | 1806-0556 | 0.122 | 0.008 | 0.004 | 0.032 | 0.230 | 0.039 | 0.019 |
| | 8/23/97 | 0605-0857 | 0.172 | 0.013 | 0.004 | 0.020 | 0.255 | 0.051 | 0.027 |
| | 8/23/97 | 0900-1156 | 0.020 | 0.004 | 0.001 | 0.005 | 0.032 | 0.025 | 0.015 |
| | 8/23/97 | 1202-1458 | 0.002 | 0.016 | 0.0003 | 0.001 | 0.010 | 0.013 | 0.009 |
| | 8/23/97 | 1500-1800 | 0.001 | 0.001 | 0.0004 | 0.0004 | 0.006 | -* | 0.004 |

*Not detected.

Azusa Intensive 2 (Aug. 4-6, 1997)




Figure 7. Biogenic hydrocarbons at Azusa during Intensive 2, Aug. 4-6, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed (others = camphene + sabinene + β -pinene + cineole + camphor) are shown.

Azusa Intensive 3 (Aug. 22-23, 1997)




Figure 8. Biogenic hydrocarbons at Azusa during Intensive 3, Aug. 22-23, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed (others = camphene + sabinene + cineole + camphor) are shown.

Azusa Intensive 4 (Sept. 4-7, 1997)




Figure 9. Biogenic hydrocarbons at Azusa during Intensive 4, Sept. 4-7, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed (others = camphene + sabinene + β -pinene + cineole + camphor) are shown.

Azusa Intensive 5 (Sept. 28-29, 1997)




Figure 10. Biogenic hydrocarbons at Azusa during Intensive 5, Sept. 28-29, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed (others = camphene + sabinene + β -pinene + cineole + camphor) are shown.

Azusa Intensive 6 (Oct. 3-4, 1997)




Figure 11. Biogenic hydrocarbons at Azusa during Intensive 6, Oct. 3-4, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed (others = camphene + sabinene + β -pinene + cineole + camphor) are shown.

Los Angeles North Main St. Intensive 3 (Aug. 22-23, 1997)




Figure 12. Biogenic hydrocarbons at Los Angeles, North Main St. during Intensive 3, Aug. 22-23, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed (others = camphene + sabinene + β -pinene + cineole + camphor) are shown.




Figure 13. Biogenic hydrocarbons at Banning during Intensive 4, Sept. 5-6, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed (others = camphene, sabinene + β -pinene + cineole + camphor) are shown.

Pine Mountain Intensive 4 (Sept. 4-7, 1997)




Figure 14. Biogenic hydrocarbons at Pine Mountain during Intensive 4, Sept. 4-7, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed (others = camphene + sabinene + β -pinene + cineole + camphor) are shown.

Mount Baldy Intensive 5 (Sept. 28-29, 1997)




Figure 15. Biogenic hydrocarbons at Mount Baldy during Intensive 5, Sept. 28-29, 1997. Ambient concentrations of isoprene, α -pinene, limonene and the sum of the remaining monoterpenes analyzed (others = camphene + sabinene + β -pinene + cineole + camphor) are shown.

5. Discussion

5.1 Biogenic Hydrocarbon Emissions

Elucidation of the mechanisms by which plants emit BHC has recently received increased attention (Turlings and Tumlinson, 1992; Monson et al., 1995; Nemecek-Marshall et al., 1995; Wildermuth and Fall, 1996; Sharkey, 1996; Lerdau et al., 1997). Isoprene is formed in plants via the mevalonic acid pathway (Sharkey et al., 1991) and its emission rate is species-specific and related to the photosynthetic rate (Loreto and Sharkey, 1993; Monson et al., 1995; Wildermuth and Fall, 1996; Sharkey, 1996; Lerdau et al., 1997). It has been proposed that the production of isoprene aids in thermal protection of plant leaves (Sharkey, 1996; Lerdau et al., 1997), but questions remain regarding why some plants, even within the same genus (e.g., oak trees) contain isoprene-emitting and nonemitting species (Fall, 1999). Isoprene does not build-up in plant tissues; its production represents a loss of fixed carbon from the plant and isoprene emission rates decline immediately when photosynthetic light is removed (Monson et al., 1991). The isoprene emission rate of a plant is a function of light intensity and temperature (Lerdau et al., 1997; Tingey et al., 1979; Guenther et al., 1991; 1993). Generally, the isoprene emission rate increases with light intensity up to a photosynthetically active radiation intensity of $\sim 1000 \mu\text{mol m}^{-2} \text{s}^{-1}$, above which the emission rate becomes constant, and increases with temperature up to a value of $\sim 40^\circ\text{C}$, above which the emission rate is reported to decrease rapidly due to denaturing of the biosynthetic enzymes (Lerdau et al., 1997; Tingey et al., 1979; Guenther et al., 1991; 1993).

Monoterpenes serve a variety of functions in plants, including a protective antiherbivore role, indirect defense by attraction of enemies of herbivores, attraction of pollinators, and allelopathic effects on competing plants (Lerdau et al., 1997; Fall, 1999). Most monoterpenoid-emitting plant species have a monoterpene reservoir pool and emissions are controlled by the temperature and the resulting leaf-air flux (Guenther et al., 1991; 1993; Tingey et al., 1980; Juuti et al., 1990). The temperature dependence of the emission rate is similar to the temperature dependence of the vapor pressure of the

monoterpene, increasing by a factor of ~2.5 for a 10°C temperature increase (Guenther et al., 1991; 1993; Tingey et al., 1980; Juuti et al., 1990). Monoterpene emissions from plants have generally not been found to be light dependent, with the most notable exception being *Quercus ilex* L., the Mediterranean Holm Oak (Staudt and Seufert, 1995; Kesselmeier et al., 1996; Loreto et al., 1996). Therefore, unlike the isoprene emissions, monoterpene emissions would be expected to continue throughout the night, however, generally at reduced levels, reflecting decreased nighttime temperatures.

5.2 Isoprene and Its Reaction Products

The highest isoprene mixing ratios were found in samples from the two mountain sites, Pine Mtn. and Mt. Baldy. Both were late afternoon/early evening samples with sampling times of 1700-2000. The highest value observed at Pine Mtn. was 2.2 ppbv on August 4, and the maximum isoprene value for Mt. Baldy was 2.3 ppbv observed on September 28. At Azusa the highest mixing ratios for isoprene were in the range 0.5 – 0.8 ppbv during all sampling intensives, except the final one on October 3-4, when the highest values were only about 0.2 ppbv. Thus, the highest values at the mountain sites were three to four times higher than the maximum mixing ratios observed at the urban valley site. The maximum isoprene values at the Banning site in the eastern portion of the air basin were 0.2 – 0.3 ppbv, consistently lower than at the other sites.

Consistent with light-dependent biogenic sources, the lowest values for isoprene were usually measured during nighttime. The diurnal variation of isoprene at the different sites generally followed a similar pattern: higher mixing ratios during the daylight hours, lower during nighttime. However, as may be seen from Figures 2, 4, 5 and 6, the difference between day and night was most apparent at the mountain sites, where the daytime values ranged between 0.4 – 2.3 ppbv and the nighttime values did not exceed 0.1 ppbv. At Azusa and Banning the day-night variation was much less pronounced and sampling at the Los Angeles North Main site gave low isoprene concentrations with little decrease during the single night sampled (see Figure 3).

5.2.1 Methacrolein and Methyl vinyl ketone

As will be discussed in more detail below, the hydroxyl (OH) radical-initiated reaction of isoprene is the major daytime loss process for isoprene and leads to an estimated atmospheric lifetime (an atmospheric lifetime is the time required for the concentration of the compound to decrease by reaction to $1/e$ of the amount initially present in the atmosphere) for isoprene of 1.4 hr (Atkinson and Arey, 1998). Methyl vinyl ketone and methacrolein are products of this rapid isoprene photooxidation reaction with yields of 0.32 and 0.22, respectively (Atkinson, 1997). Because isoprene degrades so quickly in the atmosphere, an accurate accounting of its importance in O_3 formation requires ambient measurements of its reaction products, MACR and MVK, as well as of isoprene.

For methacrolein and methyl vinyl ketone the highest mixing ratios during the SCOS97 intensive sampling periods were observed at Azusa. On August 22-23 for the 1800-0300 sampling interval the methacrolein mixing ratio was 3.3 ppbv, and on September 28 between 0300-0600 the methyl vinyl ketone mixing ratio was 2.3 ppbv. At Banning on August 22-23 at 1800-0300 mixing ratios for both MACR and MVK were high, being 2.4 ppbv and 2.9 ppbv, respectively. At Pine Mtn. and Mt. Baldy the highest values for MACR were measured in the afternoon/early evening reaching 1.1 ppbv and 0.5 ppbv, respectively, and the highest values for MVK were 1.6 and 1.0 ppbv, respectively.

At the mountain sites during the daytime, the isoprene values generally exceeded the MACR and MVK concentrations, while during nighttime the isoprene values were <0.1 ppbv and the combined MACR and MVK levels reached 1.7 ppbv. The isoprene, MACR and MVK concentrations at the mountain sites are shown in Figures 16 through 19. Note that the isoprene begins to increase early in the morning (0600-0900) with increases in MACR and MVK generally lagging behind temporally. The highest MACR and MVK levels observed at the mountain sites occurred at Pine Mtn. during Intensives 2 and 4 (see Figures 16 and 17) when the MVK + MACR reached 2.5 ppbv. The ratio MVK/MACR during the daytime (0600-2000 hr) was relatively constant at the mountain

Pine Mountain Intensive 2 (Aug. 4-6, 1997)




Figure 16. Isoprene, methacrolein and methyl vinyl ketone values at Pine Mtn. during Intensive 2. Aug. 4-6, 1997.

Pine Mountain Intensive 4 (Sept. 4-7, 1997)




Figure 17. Isoprene, methacrolein and methyl vinyl ketone values at Pine Mtn. during Intensive 4, Sept. 4-7, 1997.




Figure 18. Isoprene, methacrolein and methyl vinyl ketone at Mt. Baldy during Intensive 5, Sept. 28-29, 1997.

Mt. Baldy Intensive 6 (Oct. 3-4, 1997)




Figure 19. Isoprene, methacrolein and methyl vinyl ketone at Mt. Baldy during Intensive 6, Oct. 3-4, 1997.

sites, with average daytime values for this ratio ranging from 1.6 to 2.2. A ratio of 1.45 for MVK/MACR would be expected from the OH radical-initiated reaction of isoprene if negligible further reaction of MVK and MACR occurred, while at steady-state this ratio would be expected to increase to 2.4 (Atkinson, 1997). Thus the daytime MVK/MACR ratio at the mountain sites is consistent with the source of MACR and MVK being photooxidation of isoprene (Montzka et al. 1993, 1995; Biesenthal et al., 1997).

In contrast to the mountain sites, at Azusa, Banning, and LA North Main the MACR and MVK levels generally exceeded those of isoprene. The isoprene, MACR and MVK values for the Sept. 4-7 intensive are shown in Figures 20 and 21 for measurements taken at Azusa and Banning, respectively. The ratios of MVK/MACR were generally similar at Azusa during Intensives 2 and 4, but unusual ratios occurred during Intensives 3,5 and 6. For example, high MVK values at Azusa producing high MVK/MACR ratios were observed on Sept. 28 and Oct. 4 both for the 0300-0600 sampling period. These data suggest that a source other than the photooxidation of isoprene may exist for some of the MVK (and possibly MACR) observed at Azusa, at least during certain sampling periods. Biesenthal and co-investigators suggested that a significant non-photochemical source, possibly automotive in origin, of MVK and MACR was present in an urban area of British Columbia (Biesenthal et al., 1997). The MVK/MACR ratios observed in the early mornings of Sept. 28 and Oct. 4, however, are much higher than ratios observed in emissions from several types of vehicles (Jonsson et al., 1985) and also higher than the ratio observed in Toronto, Canada in the winter, where the lack of isoprene emissions from plant species and a correlation of the MVK and MACR with CO implicated automotive sources (Biesenthal and Shepson, 1997). If automobiles are a source of MVK and MACR they can be expected to make a contribution to the levels of these species measured at Azusa, a heavily traffic-impacted site. Further interpretation of the MVK and MACR data may be possible when the anthropogenic VOC data and gaseous co-pollutant data (CO, NO, O₃, etc.) for Azusa and Banning become available.

Asuza Intensive 4 (Sept. 4-7, 1997)




Figure 20. Isoprene, methacrolein and methyl vinyl ketone at Azusa during Intensive 4, Sept. 4-7, 1997.

Banning Intensive 4 (Sept. 5-6, 1997)




Figure 21. Isoprene, methacrolein and methyl vinyl ketone at Banning during Intensive 4, Sept. 5-6, 1997.

5.3 Monoterpenes

As noted above, most monoterpene-emitting plants have a reservoir pool of these compounds and their emissions increase with increasing ambient temperatures. Therefore, the highest monoterpene emissions from plants would be expected in the afternoon when the ambient temperatures are highest. The term monoterpenes as used here, designates not only the C₁₀H₁₆ alkenes, but also the oxygenated species such as 1,8-cineole and camphor which are emitted by vegetation (see Figure 1 for structures). As will be discussed in more detail below, the atmospheric lifetimes of C₁₀H₁₆ alkenes are generally a few hours or less. The ambient concentrations of the monoterpenes, therefore, will reflect their emission profiles, meteorology, including inversion heights and transport, as well as atmospheric reactions.

The highest monoterpene mixing ratios measured were 3.8 ppbv and 2.3 ppbv for limonene at Azusa 0600-0900 hr. on Sept. 28 and 0300-0600 hr. on Aug. 23, respectively. The average mixing ratios for limonene at Azusa ranged from 0.1 to 0.57 ppbv for Intensives 2-6. While, the average mixing ratios for α -pinene at Azusa ranged from 0.04 to 0.19, and α -pinene reached a maximum of only 0.5 ppbv, the α -pinene and limonene levels were often very comparable (see Figures 7-11), suggesting that the emission "signature" of the Azusa vegetation could have comparable levels of these monoterpenes. It seems unlikely that the very high early-morning limonene mixing ratio peaks could be the result of a biogenic emission, but rather suggest an anthropogenic source for this limonene.

For camphene, sabinene, β -pinene, cineole and camphor the mixing ratio maxima at Azusa were (in ppbv): 0.14, 0.38, 0.10, 0.10 and 0.06, respectively. Thus, the limonene maximum was nearly an order of magnitude higher than any of the other monoterpenes analyzed. As may be seen from examining Figures 7 through 11, the mixing ratios of isoprene, α -pinene, limonene and other monoterpenes will all need to be considered when determining the contribution of BHC to O₃ formation at Azusa. At the downwind Banning site (see Figure 13), isoprene and the monoterpenes are again observed in comparable levels, although isoprene is generally the highest single BHC.

At the mountain sites (see Figures 14 and 15), the BHC profiles are clearly dominated by isoprene. The mixing ratio maximum at Pine Mountain for each BHC was [compound (maximum in ppbv)]: isoprene (2.2), α -pinene (0.05), camphene (0.01), sabinene (0.03), β -pinene (0.01), limonene (0.01), cineole (0.06) and camphor (0.06). Thus, α -pinene, cineole and camphor are prominent in the monoterpene profile observed at Pine Mountain. The mixing ratio maximum at Mt. Baldy for each BHC was [compound (maximum in ppbv)]: isoprene (2.3), α -pinene (0.2), camphene (0.01), sabinene (0.02), β -pinene (0.06), limonene (0.06), cineole (0.04) and camphor (0.04). The presence of numerous coniferous trees at the Mt. Baldy site explains the slight dominance of α - and β -pinene in the monoterpene profile at this site.

5.4 Atmospheric Reactions

As noted above, many of the BHC are highly reactive in the atmosphere. Table 11 gives the rate constants for the reactions of the BHC reported on here (including the isoprene products, MACR and MVK) with the hydroxyl (OH) radical, with the nitrate (NO_3) radical, and with ozone (O_3). No kinetic data are available for camphor. Table 12 shows the calculated lifetimes for each of these BHC; including photolysis lifetimes for the carbonyls, MACR and MVK, the only compounds measured that are likely to absorb actinic radiation. To calculate these lifetimes, reasonable values for ambient OH, NO_3 and O_3 concentrations have been assumed. A 12-hr daytime average for OH of 2.0×10^6 molecules cm^{-3} was chosen based on a diurnally, seasonally, and annually averaged global tropospheric OH concentration (Prinn et al., 1995) and assuming the presence of the photolytically-formed OH only during daylight hours. Because the NO_3 radical rapidly photolyzes during the day, a 12-hr nighttime value for NO_3 of 5×10^8 molecule cm^{-3} is used (Atkinson, 1991). The lifetimes due to reaction with O_3 are calculated for ~30 ppbv of O_3 , a typical background clean troposphere value (Logan, 1985). If data become available, for example, if ambient O_3 concentration profiles are known, site- and time-specific calculations of the lifetimes can be made. It should be noted that both NO_3 and O_3 react rapidly with NO and these reactions may keep their concentrations low in urban areas where vehicles are emitting significant amounts of NO.

Table 11. Rate constants k at 298 K and 760 Torr total pressure of air for the reaction of selected biogenic hydrocarbons with the OH radical, the NO₃ radical, and O₃ (data from Atkinson, 1997 unless noted otherwise).

| Biogenic Hydrocarbon | k (cm ³ molecule ⁻¹ s ⁻¹) for reaction with | | |
|----------------------|---|--|--|
| | OH (10 ¹² x k _{OH}) | NO ₃ (10 ¹⁴ x k _{NO₃}) | O ₃ (10 ¹⁸ x k _{O₃}) |
| Isoprene | 101 | 67.8 | 12.8 |
| Methacrolein | 28.6 ^a | 0.33 ^b | 1.14 ^d |
| Methyl vinyl ketone | 20.2 ^a | <0.012 ^c | 4.56 ^d |
| α -Pinene | 53.7 | 616 | 86.6 |
| Camphene | 53 | 66 | 0.90 |
| Sabinene | 117 | 1000 | 86 |
| β -Pinene | 78.9 | 251 | 15 |
| Limonene | 171 | 1220 | 200 |
| 1,8-Cineole | 11.1 ^e | 0.017 ^e | <0.15 ^d |

^aData from Atkinson et al., 1983 and Gierczak et al., 1997.

^bData from Chew et al., 1998.

^cData from Rudich et al., 1996.

^dData from Atkinson, 1994.

^eData from Corchnoy and Atkinson, 1990.

Table 12. Calculated lifetimes^a for selected biogenic hydrocarbons with respect to photolysis, reaction with the OH radical, reaction with the NO₃ radical, and reaction with O₃.

| Biogenic Hydrocarbon | Lifetime due to | | | |
|----------------------|-----------------|------------------------------|-----------------------------|---------------------|
| | OH ^b | NO ₃ ^c | O ₃ ^d | Photolysis |
| Isoprene | 1.4 hr | 49 min | 1.3 day | |
| Methacrolein | 4.9 hr | 14 day | 15 day | ~1 day ^e |
| Methyl vinyl ketone | 6.9 hr | >385 day | 3.6 day | ~2 day ^e |
| α-Pinene | 2.6 hr | 5 min | 4.6 hr | |
| Camphene | 2.6 hr | 50 min | 18 day | |
| Sabinene | 1.2 hr | 3 min | 4.6 hr | |
| β-Pinene | 1.8 hr | 13 min | 1.1 day | |
| Limonene | 49 min | 3 min | 2.0 hr | |
| 1,8-Cineole | 12.5 hr | 270 day | >110 day | |

^aAn atmospheric lifetime is defined as the time required for the compound concentration to degrade to a value of 1/e of its initial concentration.

^bFor a 12-hr daytime average OH radical concentration of 2.0×10^6 molecule cm⁻³.

^cFor a 12-hr nighttime average NO₃ radical concentration of 5×10^8 molecule cm⁻³.

^dFor a 24-hr average O₃ concentration of 7×10^{11} molecule cm⁻³ (~30 ppbv).

^eFor overhead sun (Atkinson, 1999).

Although Table 12 gives "typical or average" lifetimes it will allow the identification of the likely most important loss processes for a given BHC. For example, isoprene loss during the daytime will be from the OH radical-initiated reaction. If NO₃ radicals are present at night, a rapid loss of isoprene can be expected. Additionally, Table 12 can be used to compare the effect of a given loss process on different BHC. For example, high O₃ levels will have the greatest effect on the lifetime of limonene, the most reactive of the BHC shown with O₃, but will have essentially no effect on the mixing ratios of 1,8-cineole.

As noted above, the isoprene concentrations at the mountain sites decreased rapidly during the nighttime. Both isoprene emissions and isoprene reaction with the OH radical will essentially stop after sunset. Potential nighttime OH sources, such as monoterpene-O₃ reactions (Atkinson et al., 1992; Paulson and Orlando, 1996) are expected to be insignificant. As noted above (see Figures 16-19), the MACR and MVK mixing ratios at the mountain sites did not decrease as much as the isoprene mixing ratios during the nighttime. Reaction of isoprene with O₃ would also produce MACR and MVK as products, in yields of 0.39 and 0.16, respectively (Atkinson, 1997), but the rate of the isoprene-O₃ reaction is not rapid enough (see Tables 11 and 12) to result in the <0.1 ppbv isoprene mixing ratios seen during the 2000-0300 sampling periods at Pine Mtn. and Mt. Baldy. For example, if the O₃ mixing ratio were 120 ppbv, the lifetime of isoprene due to reaction with O₃ would be ~7 hours. However, the rapid reaction of NO₃ with isoprene could explain the low isoprene mixing ratios during the 2000-0300 sampling periods, which then remained low until the daytime emission of isoprene began after sunrise. [It should be noted that the NO₃-isoprene reaction gives formation yields of MACR and MVK of only 0.035 each (Atkinson, 1997)]. Examining the profiles of other BHC measured at these sites provides additional evidence for the occurrence of nighttime NO₃ chemistry at the mountain sites.

Figures 22 and 23 show the mixing ratios of isoprene, α -pinene, 1,8-cineole and camphor at Pine Mountain. As may be seen from Table 12, α -pinene will react rapidly with the NO₃ radical, while 1,8-cineole will not. Although data are not available for

Pine Mtn. Intensive 2 (Aug. 4-6, 1997)




Figure 22. Isoprene, α -pinene, cineole and camphor values at Pine Mtn. during Intensive 2, Aug. 4-6, 1997.

Pine Mtn. Intensive 4 (Sept. 4-7, 1997)




Figure 23. Isoprene, α -pinene, cineole and camphor values at Pine Mtn. during Intensive 4, Sept. 4-7, 1997.

camphor, it is estimated that the reactions of camphor with NO_3 and with O_3 will be slower than those of 1,8-cineole (R. Atkinson, personal communication). Thus, the rapid decrease in isoprene and α -pinene mixing ratios during the 2000-0300 hr. sampling time, seen twice during Intensive 2 (Figure 22) and three times during Intensive 4 (Figure 23) is consistent with NO_3 radical reaction with isoprene and α -pinene. The fact that the 1,8-cineole and camphor mixing ratios, and, as mentioned earlier, the MACR and MVK mixing ratios, did not decrease to the same extent as the isoprene and α -pinene mixing ratios, rules out a meteorological explanation (such as the inversion lowering and Pine Mtn. being in the "free troposphere" during the nighttime) for the isoprene and α -pinene decreases.

Shown in Figures 24 and 25 are the isoprene, α -pinene, limonene and 1,8-cineole mixing ratios from Mt. Baldy during Intensives 5 and 6. Again, the isoprene mixing ratio decreases dramatically between the 1700-2000 hr. sample and the nighttime samples. At Mt. Baldy during Intensive 5, the α -pinene and limonene mixing ratios decreased more than the 1,8-cineole mixing ratio, but not as much as that of isoprene (Figure 24). While the emissions of isoprene will stop at night, the monoterpenes will continue to be emitted, and fresh emissions of α -pinene and limonene at Mt. Baldy may explain why the mixing ratios of these monoterpenes measured at night did not decrease proportionally as much as those of isoprene. As mentioned previously, the Mt. Baldy site had significantly more coniferous trees than the Pine Mtn. site, and this is reflected in the higher α -pinene/isoprene ratio at Mt. Baldy during Intensive 5 relative to Pine Mtn.

6. Conclusions and Implications

Detectable levels of the biogenic hydrocarbons, isoprene, α -pinene, camphene, sabinene, β -pinene, limonene, 1,8-cineole and camphor were observed in downtown Los Angeles (L.A. North Main St.), at a mid-basin receptor site (Azusa), at mountain locations (Pine Mtn. and Mt. Baldy) and at a far downwind receptor site (Banning) during the SCOS97 study. Isoprene concentrations were highest at the two mountain sites, Pine

Mt. Baldy Intensive 5 (Sept. 28-29, 1997)




Figure 24. Isoprene, α -pinene, limonene and cineole values at Mt. Baldy during Intensive 5, Sept 28-29, 1997.

Mt. Baldy Intensive 6 (Oct. 3-4, 1997)




Figure 25. Isoprene, α -pinene, limonene and cineole values at Mt. Baldy during Intensive 6, Oct. 3-4, 1997.

Mtn. and Mt. Baldy, and showed very pronounced diurnal profiles consistent with light-dependent biogenic emission sources. The maximum isoprene concentrations observed at these sites were 2.2 and 2.3 ppbv, respectively, both during the 1700-2000 hr. sampling period. At the mountain site during the daytime, the isoprene values generally exceeded the MACR and MVK concentrations and the MVK/MACR ratios were consistent with their source being photooxidation of isoprene. The rate constants for atmospheric reactions with the OH radical, the NO₃ radical and O₃ of the BHC measured were examined to explore the effect of atmospheric chemistry on the diurnal profiles measured. The rapid decreases at the Pine Mountain site in the isoprene and α-pinene mixing ratios between the 1700-2000 hr. sampling period and the nighttime sampling period (2000-0300) which were not mirrored in mixing ratio changes for 1,8-cineole and camphor suggest that NO₃ was present during the night at this mountain site causing rapid reaction of isoprene and α-pinene.

In contrast to the mountain sites, at Azusa, Banning, and LA North Main the MACR and MVK levels generally exceeded those of isoprene, reaching combined MACR + MVK values of 4.9, 5.3 and 2.2 ppbv, respectively. The isoprene concentrations were generally highest during daytime at these sites, but did not decrease at night as much as at the mountain sites. Occasional, unusual MVK/MACR ratios at Azusa and Banning suggest potential anthropogenic sources for these compounds, and the anthropogenic VOC data from these sites should facilitate further interpretation of the data for isoprene and its atmospheric reaction products.

The monoterpene observed in highest concentration was limonene, which reached 3.8 ppbv at Azusa during the early morning (0600-0900) of Sept. 28. During the early morning hours (0300-0900) on Aug. 22, high limonene mixing ratios were also observed at Azusa. The dominance of this single monoterpene, during time periods when low ambient temperatures would not result in enhanced biogenic emissions, suggests that limonene at Azusa has an occasional strong anthropogenic source, as well as biogenic sources. During the daytime, the mountain sites were characterized by isoprene as the dominant BHC. In contrast, at Azusa and Banning, the monoterpenes made a significant contribution to the daytime BHC total.

7. References

- Atkinson, R. (1994) Gas-phase tropospheric chemistry of organic compounds. *J. Phys. Chem. Ref. Data, Monograph No. 2*, 216 pp.
- Atkinson, R. (1997) Gas-phase tropospheric chemistry of volatile organic compounds:1. Alkanes and alkenes. *J. Phys. Chem. Ref. Data*, 26, 215-290.
- Atkinson, R. (1999) Atmospheric Chemistry of VOCs and NO_x. *Atmos. Environ.*, *in press*.
- Atkinson, R. and J. Arey (1998) Atmospheric chemistry of biogenic organic compounds. *Accounts of Chemical Research*, 31, 574-583.
- Atkinson, R., S. M. Aschmann, and J. N. Pitts, Jr. (1983) Kinetics of the gas-phase reactions of OH radicals with a series of α,β -unsaturated carbonyls at 299 ± 2 K. *Int. J. Chem. Kinet.*, 15, 75-81.
- Atkinson, R., S. M. Aschmann, J. Arey, and B. Shorees (1992) Formation of OH radicals in the gas phase reactions of O₃ with a series of terpenes. *J. Geophys. Res.*, 97, 6065-6073.
- Benjamin, M. T., M. Sudol, D. Vorsatz, and A. M. Winer (1997) A spacially and temporally resolved biogenic hydrocarbon emissions inventory for the California south coast air basin. *Atmos. Environ.*, 31, 3087-3100.
- Biesenthal, T. A. and P. B. Shepson (1997) Observations of anthropogenic inputs of the isoprene oxidation products methyl vinyl ketone and methacrolein to the atmosphere. *Geophys. Res. Lett.*, 24, 1375-1378.
- Biesenthal, T. A., Q. Wu, P. B. Shepson, H. A. Wiebe, K. G. Anlauf, and G. I. Mackay (1997) A study of relationships between isoprene, its oxidation products, and ozone, in the lower Fraser Valley, BC. *Atmos. Environ.*, 31, 2049-2058.
- Carter, W. P. L. (1994) Development of ozone reactivity scales for volatile organic compounds. *J. Air & Waste manage. Assoc.*, 44, 881-899.
- Chameides, W. L., R. W. Lindsay J. Richardson, and C. S. Kiang (1988) The role of biogenic hydrocarbons in urban photochemical smog: Atlanta as a case study. *Science*, 241, 1473-1475.
- Chew, A.A., R. Atkinson, and S. M. Aschmann (1998) Kinetics of the gas-phase reaction of NO₃ radicals with a series of alcohols, glycol ethers, ethers and chloroalkenes. *J. Chem. Soc. Faraday Trans.*, 94, 1083-1089.

Corchnoy, S. B. and R. Atkinson (1990) Kinetics of the gas-phase reactions of OH and NO₃ radicals with 2-carene, 1,8-cineole, p-cymene, and terpinolene. *Environ. Sci. Technol.*, 24, 1497-1502.

Fall, R. (1999) Biogenic emissions of volatile organic compounds from higher plants. IN: *Reactive Hydrocarbons in the Atmosphere*, Academic Press, pp. 41-96.

Gierczak, T., J. B. Burkholder, R. K. Talukdar, A. Mellouki, S. B. Barone, and A. R. Ravishankara (1997) Atmospheric fate of methyl vinyl ketone and methacrolein. *J. Photochem. and Photobiol. A: Chemistry*, 110, 1-10.

Guenther, A. B., R. K. Monson, and R. Fall (1991) Isoprene and monoterpene emission rate variability: observations with eucalyptus and emission rate algorithm development. *J. Geophys. Res.*, 96, 10799-10808.

Guenther, A. B., P. R. Zimmerman, and P. C. Harley (1993) Isoprene and monoterpene emission rate variability: model evaluations and sensitivity analyses. *J. Geophys. Res.*, 98, 12609-12617.

Guenther, A., P. Zimmerman, and M. Wildermuth (1994) Natural volatile organic compound emission rate estimates for U.S. woodland landscapes. *Atmos. Environ.*, 28, 1197-1210.

Guenther, A., C. Hewitt, D. Erickson, R. Fall, C. Geron, T. Graedel, P. Harley, L. Klinger, M. Lerdau, W. McKay, T. Pierce, B. Scholes, R. Steinbrecher, R. Tallamraju, J. Taylor, and P. Zimmerman (1995) A global model of natural volatile organic compound emissions. *J. Geophys. Res.*, 100, 8873-8892.

Helwig, D. (1997) Ozone removal techniques in the sampling of atmospheric volatile organic trace gases. *Atmos. Environ.*, 31, 3635-3651.

Hoffman, T. (1994) Adsorptive preconcentration technique including oxidant scavenging for the measurement of reactive natural hydrocarbons in ambient air. *Fres. J. Anal. Chem.*, 348, 41541-41547.

Jonsson, A., K. A. Persson, and V. Grigoriadis (1985) Measurements of some low molecular-weight oxygenated, aromatic, and chlorinated hydrocarbons in ambient air and in vehicle emissions. *Environ. Int.*, 11, 383-392.

Juttner, F. (1988) A cryotrap technique for the quantitation of monoterpenes in humid and ozone-rich forest air. *J. Chromatogr.*, 442, 157-163.

Juuti, S., J. Arey, and R. Atkinson (1990) Monoterpene emission rate measurements from a Monterey pine. *J. Geophys. Res.*, 95, 7515-7519.

Kesselmeier, J., L. Schafer, P. Ciccioli, E. Brancaleoni, A. Cecinato, M. Frattoni, P. Foster, V. Jacob, J. Denis, J. L. Fugit, L. Dutaur, and L. Torres (1996) Emission of monoterpenes and isoprene from a Mediterranean oak species *Quercus ilex* L. measured within the BEMA (Biogenic emissions in the Mediterranean area) project. *Atmos. Environ.*, 30, 1841-1850.

Lerdau, M., A. Guenther, and R. Monson (1997) Plant production and emission of volatile organic compounds. *BioScience*, 47, 373-383.

Loreto, F. and T. D. Sharkey (1993) On the relationship between isoprene emission and photosynthetic metabolites under different environmental conditions. *Planta*, 189, 420-424.

Loreto, F., P. Ciccioli, A. Cecinato, E. Brancaleoni, M. Frattoni, C. Fabozzi, and D. Tricoli (1996) Evidence of the photosynthetic origin of monoterpenes emitted by *Quercus ilex* L. leaves by ^{13}C labeling. *Plant Physiol.*, 110, 1317-1322.

Monson, R. K., A. J. Hills, P. R. Zimmerman, and R. R. Fall (1991) Studies of the relationship between isoprene emission rate and CO_2 or photon-flux density using a real-time isoprene analyzer. *Plant, Cell and Environment*, 14, 517-523.

Monson, R. K., M. T. Lerdau, T. D. Sharkey, D. S. Schimel, and R. Fall (1995) Biological aspects of constructing volatile organic compound emission inventories. *Atmos. Environ.*, 29, 2989-3002.

Montzka, S. A., M. Trainer, P. D. Goldan, W. C. Kuster, and F. C. Fehsenfeld (1993) Isoprene and its oxidation products, methyl vinyl ketone and methacrolein, in the rural troposphere. *J. Geophys. Res.*, 98, 1101-1111.

Montzka, S. A., M. Trainer, W. M. Angevine, and F. C. Fehsenfeld (1995) Measurements of 3-methylfuran, methyl vinyl ketone, and methacrolein at a rural forested site in the southeastern United States. *J. Geophys. Res.*, 100, 11393-11401.

Nemecek-Marshall, M., R. C. MacDonald, J. J. Franzen, C. L. Wojciechowski, and R. Fall (1995) Methanol emission from leaves. *Plant Physiol.*, 108, 1359-1368.

Paulson, S. E. and J. J. Orlando (1996) The reactions of ozone with alkenes: an important source of HO_x in the boundary layer. *Geophys. Res. Lett.*, 23, 3727-3730.

Rudich, Y., R. K. Talukdar, R. W. Fox, and A. R. Ravishankara (1996) Rate coefficients for reactions of NO_3 with a few olefins and oxygenated olefins. *J. Phys. Chem.*, 100, 5374-5381.

Sharkey, T. D., F. Loreto, and C. F. Delwiche (1991) The biochemistry of isoprene emission from leaves during photosynthesis. IN: *Trace Gas Emissions by Plants*, T. D. Sharkey, E. A. Holland, and H. A. Mooney, eds., Academic Press pp. 153-184.

Sharkey, T. D. (1996) Emission of low molecular mass hydrocarbons from plants. *Trends in Plant Science*, 1, 78-82.

Staudt, M. and G. Seufert (1995) Light-dependent emission of monoterpenes by Holm Oak (*Quercus ilex* L.). *Naturwissenschaften*, 82, 89-92.

Tingey, D. T., M. Manning, L. C. Grothaus, and W. F. Burns (1979) The influence of light and temperature on isoprene emission rates from live oak. *Plant Physiol.*, 47, 112-118.

Tingey, D. T., M. Manning, L. C. Grothaus, and W. F. Burns (1980) Influence of light and temperature on monoterpene emission rates from slash pine. *Plant Physiol.*, 65, 797-801.

Trainer, M., E. Y. Hsie, S. A. McKeen, R. Tallamraju, D. D. Parrish, F. C. Fehsenfeld, and S. C. Liu (1987) Impact of natural hydrocarbons on hydroxyl and peroxy radicals at a remote site. *J. Geophys. Res.*, 92, 11879-11894.

Turlings, T. C. J. and J. H. Tumlinson (1992) Systemic release of chemical signals by herbivore-injured corn. *Proc. Natl. Acad. Sci. U.S.A.*, 89, 8399-8402.

Wildermuth, M. C. and R. Fall (1996) Light-dependent isoprene emission. *Plant Physiol.*, 112, 171-182.

WMO (1995) World Meteorological Organization, *Scientific Assessment of Ozone Depletion: 1994*, Global Ozone Research and Monitoring Project - Report No. 37, Chapter 2, Geneva, Switzerland.

Table A-1.1. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Azusa.

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|----------------------------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|----------------------------|
| AZUSA Aug 4-6, 1997 | | | | | | | | |
| 8/4/97 | 0600-0900 | 0.23 | 0.23 | 0.11 | 0.09 | 0.41 | 0.46 | |
| 8/4/97 | 0600-0900 | 0.23 | | 0.08 | | 0.52 | | |
| 8/4/97 | 0907-1206 | 0.51 | 0.52 | 0.48 | 0.47 | 0.76 | 0.77 | |
| 8/4/97 | 0907-1206 | 0.53 | | 0.46 | | 0.78 | | |
| 8/4/97 | 1306-1603 | 0.46 | 0.46 | 0.65 | 0.51 | 1.06 | 0.92 | |
| 8/4/97 | 1306-1603 | 0.47 | | 0.36 | | 0.78 | | |
| 8/4/97 | 1700-2000 | 0.74 | 0.75 | 0.56 | 0.71 | 0.79 | 1.06 | |
| 8/4/97 | 1700-2000 | 0.76 | | 0.86 | | 1.33 | | |
| 8/4/97 | 2006-0300 | 0.14 | 0.16 | 0.33 | 0.34 | 0.50 | 0.59 | |
| 8/4/97 | 2006-0300 | 0.18 | | 0.34 | | 0.69 | | |
| 8/5/97 | 0307-0600 | 0.23 | 0.24 | 1.01 | 1.01 | 0.57 | 0.58 | |
| 8/5/97 | 0307-0600 | 0.24 | | 1.02 | | 0.58 | | |
| 8/5/97 | 0618-0900 | 0.46 | 0.47 | 0.72 | 0.82 | 1.46 | 1.73 | |
| 8/5/97 | 0618-0900 | 0.48 | | 0.92 | | 2.00 | | |
| 8/5/97 | 0907-1202 | 0.63 | 0.66 | 0.72 | 0.74 | 1.55 | 1.61 | |
| 8/5/97 | 0907-1202 | 0.70 | | 0.75 | | 1.68 | | |
| 8/5/97 | 1300-1600 | 0.59 | 0.61 | 0.61 | 0.70 | 1.29 | 1.49 | |
| 8/5/97 | 1300-1600 | 0.62 | | 0.78 | | 1.70 | | |
| 8/5/97 | 1701-2000 | 0.74 | 0.73 | 0.45 | 0.46 | 0.70 | 0.72 | |
| 8/5/97 | 1701-2000 | 0.72 | | 0.48 | | 0.75 | | |
| 8/5- 8/6/97 | 2005-0300 | 0.47 | 0.44 | 0.42 | 0.41 | 1.22 | 1.20 | |
| 8/5-8/6/97 | 2005-0300 | 0.41 | | 0.41 | | 1.19 | | |
| 8/6/97 | 0306-0600 | 0.23 | 0.23 | 0.60 | 0.62 | 0.90 | 0.90 | |
| 8/6/97 | 0306-0600 | 0.23 | | 0.64 | | 0.89 | | |
| 8/6/97 | 0616-0900 | 0.37 | 0.37 | 0.65 | 0.60 | 1.26 | 1.24 | |
| 8/6/97 | 0616-0900 | 0.38 | | 0.55 | | 1.23 | | |

Table A-1.1. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Azusa
(continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|---------------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|----------------------------|
| 8/6/97 | 0906-1200 | 0.50 | 0.51 | 0.37 | 0.35 | 0.74 | 0.75 | |
| 8/6/97 | 0906-1200 | 0.52 | | 0.33 | | 0.75 | | |
| 8/6/97 | 1300-1600 | 0.51 | 0.51 | 0.45 | 0.44 | 0.83 | 0.81 | |
| 8/6/97 | 1300-1600 | 0.51 | | 0.43 | | 0.79 | | |
| 8/6/97 | 1700-2000 | 0.40 | | 0.11 | 0.15 | 0.17 | 0.23 | |
| 8/6/97 | 1700-2000 | 0.40 | 0.18 | | | 0.29 | | |
| AZUSA | | | | | | | | |
| 8/22/97 | 0318-0600 | 0.24 | 0.24* | 0.29 | 0.29* | 0.53 | 0.53* | |
| 8/22/97 | 0318-0600 | 0.67 | | 1.15 | | 2.04 | | |
| 8/22/97 | 0610-0900 | 0.02 | 0.21* | 0.09 | 0.45* | 0.08 | 0.77* | |
| 8/22/97 | 0610-0900 | 0.21 | | 0.45 | | 0.77 | | |
| 8/22/97 | 0905-1157 | 0.39 | 0.41 | 0.34 | 0.33 | 0.66 | 0.69 | |
| 8/22/97 | 0905-1157 | 0.44 | | 0.32 | | 0.71 | | |
| 8/22/97 | 1210-1500 | 0.61 | 0.52* | 1.33 | 0.44* | 1.50 | 0.78* | |
| 8/22/97 | 1210-1500 | 0.52 | | 0.44 | | 0.78 | | |
| 8/22/97 | 1504-1800 | 0.47 | 0.45 | 0.79 | 0.69 | 1.17 | 1.07 | |
| 8/22/97 | 1504-1800 | 0.44 | | 0.59 | | 0.97 | | |
| 8/22-23/97 | 1815-0300 | 0.28 | 0.30 | 2.82 | 3.34 | 1.01 | 1.52 | |
| 8/22-23/97 | 1815-0300 | 0.32 | | 3.87 | | 2.03 | | |
| 8/23/97 | 0304-0600 | 0.27 | 0.25 | 0.45 | 0.55 | 1.16 | 0.98 | |
| 8/23/97 | 0304-0600 | 0.22 | | 0.65 | | 0.80 | | |
| 8/23/97 | 0615-0900 | 0.26 | 0.26* | 0.70 | 0.70* | 0.94 | 0.94* | |
| 8/23/97 | 0615-0900 | - | | - | - | - | - | |

(Mass detector stopped during analysis, sample destroyed)

Table A-1.1. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Azusa (continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|---------------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|---|
| 8/23/97 | 0903-1200 | 0.53 | 0.51 | 0.57 | 0.53 | 0.61 | 0.74 | |
| 8/23/97 | 0903-1200 | 0.48 | | 0.49 | | 0.88 | | |
| 8/23/97 | 1208-1500 | 0.34 | 0.31 | 0.56 | 0.49 | 0.72 | 0.82 | |
| 8/23/97 | 1208-1500 | 0.29 | | 0.41 | | 0.92 | | |
| 8/23/97 | 1505-1800 | 0.37 | 0.37* | 0.46 | 0.46* | 0.70 | 0.70* | |
| 8/23/97 | 1505-1800 | 0.42 | | 0.55 | | 0.98 | | Sampling tube contents transferred from broken tube |
| AZUSA | | | | | | | | |
| Sep 4-7, 1997 | | | | | | | | |
| 9/4/97 | 0300-0600 | 0.27 | 0.28 | 0.88 | 0.89 | 1.43 | 1.29 | |
| 9/4/97 | 0300-0600 | 0.28 | | 0.89 | | 1.15 | | |
| 9/4/97 | 0606-0900 | 0.22 | 0.22 | 0.55 | 0.55 | 0.63 | 0.56 | |
| 9/4/97 | 0606-0900 | 0.22 | | 0.55 | | 0.50 | | |
| 9/4/97 | 0904-1200 | 0.58 | 0.57 | 0.49 | 0.46 | 1.77 | 1.54 | |
| 9/4/97 | 0904-1200 | 0.56 | | 0.44 | | 1.31 | | |
| 9/4/97 | 1300-1600 | 0.35 | 0.35 | 0.38 | 0.62 | 0.75 | 1.00 | |
| 9/4/97 | 1300-1600 | 0.35 | | 0.86 | | 1.25 | | |
| 9/4/97 | 1700-2000 | 0.59 | 0.61 | 0.34 | 0.40 | 0.63 | 0.50 | |
| 9/4/97 | 1700-2000 | 0.63 | | 0.47 | | 0.36 | | |
| 9/4/97 | 2014-2400 | 0.14 | 0.13 | 0.29 | 0.26 | 0.28 | 0.25 | |
| 9/4/97 | 2014-2400 | 0.12 | | 0.23 | | 0.22 | | |
| 9/5/97 | 0006-0300 | 0.07 | 0.06 | 0.15 | 0.17 | 0.32 | 0.26 | |
| 9/5/97 | 0006-0300 | 0.06 | | 0.20 | | 0.20 | | |
| 9/5/97 | 0305-0600 | 0.06 | 0.06 | 0.15 | 0.16 | 0.30 | 0.25 | |
| 9/5/97 | 0305-0600 | 0.06 | | 0.16 | | 0.19 | | |
| 9/5/97 | 0615-0900 | - | 0.17* | - | 0.48* | - | 0.38* | High toluene, analysis in the SCAN mode |
| 9/5/97 | 0615-0900 | 0.17 | | 0.48 | | 0.38 | | |
| 9/5/97 | 0903-1200 | 0.40 | 0.43 | 0.57 | 0.57* | 0.98 | 0.98* | Analyzed much later than other samples |
| 9/5/97 | 0903-1200 | 0.46 | | 0.05 | | 0.38 | | |

Table A-1.1. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Azusa (continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|--------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|----------------------------|
| 9/5/97 | 1300-1600 | 0.32 | 0.31 | 0.41 | 0.39 | 1.08 | 1.01 | |
| 9/5/97 | 1300-1600 | 0.30 | | 0.37 | | 0.94 | | |
| 9/5/97 | 1700-2000 | 0.36 | 0.37 | 0.27 | 0.32 | 0.33 | 0.41 | |
| 9/5/97 | 1700-2000 | 0.38 | | 0.37 | | 0.48 | | |
| 9/5/97 | 2006-2400 | 0.19 | 0.18 | 0.37 | 0.58 | 0.42 | 0.57 | |
| 9/5/97 | 2006-2400 | 0.18 | | 0.78 | | 0.73 | | |
| 9/6/97 | 0008-0300 | 0.11 | 0.11 | 0.22 | 0.22 | 0.33 | 0.28 | |
| 9/6/97 | 0008-0300 | 0.10 | | 0.23 | | 0.24 | | |
| 9/6/97 | 0306-0600 | 0.09 | 0.09 | 0.23 | 0.30 | 0.36 | 0.36 | |
| 9/6/97 | 0306-0600 | 0.10 | | 0.37 | | 0.37 | | |
| 9/6/97 | 0615-0900 | 0.14 | 0.15 | 0.30 | 0.33 | 0.27 | 0.27 | |
| 9/6/97 | 0615-0900 | 0.15 | | 0.35 | | 0.27 | | |
| 9/6/97 | 0903-1200 | 0.34 | 0.35 | 0.48 | 0.50 | 0.54 | 0.65 | |
| 9/6/97 | 0903-1200 | 0.35 | | 0.53 | | 0.76 | | |
| 9/6/97 | 1300-1600 | 0.30 | 0.29 | 0.43 | 0.44 | 0.89 | 0.88 | |
| 9/6/97 | 1300-1600 | 0.28 | | 0.45 | | 0.86 | | |
| 9/6/97 | 1700-1959 | 0.29 | 0.29 | 0.31 | 0.32 | 0.44 | 0.50 | |
| 9/6/97 | 1700-1959 | 0.30 | | 0.32 | | 0.56 | | |
| 9/6/97 | 2010-2400 | 0.09 | 0.08 | 0.18 | 0.16 | 0.17 | 0.18 | |
| 9/6/97 | 2010-2400 | 0.08 | | 0.13 | | 0.18 | | |
| 9/7/97 | 0005-0300 | 0.10 | 0.11 | 0.20 | 0.21 | 0.22 | 0.23 | |
| 9/7/97 | 0005-0300 | 0.11 | | 0.22 | | 0.23 | | |
| 9/7/97 | 0306-0600 | 0.11 | 0.11 | 0.29 | 0.30 | 0.47 | 0.52 | |
| 9/7/97 | 0306-0600 | 0.12 | | 0.31 | | 0.57 | | |
| 9/7/97 | 0604-0900 | 0.12 | 0.13 | 0.31 | 0.35 | 0.37 | 0.31 | |
| 9/7/97 | 0604-0900 | 0.13 | | 0.39 | | 0.25 | | |

Table A-1.1. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Azusa (continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|--------------|------------------------------|------------------|----------------------|--------------|------------------|-------------|-----------------|-------------------------------------|
| AZUSA | | | | | | | | |
| 9/28/97 | Sep 28-29, 1997 0301-0600 | 0.16 | 0.15 | 0.32 | 0.28 | 2.30 | 2.31 | |
| 9/28/97 | 0301-0600 | 0.14 | | 0.25 | | 2.33 | | |
| 9/28/97 | 0609-0900 | 0.23 | 0.23 | - | 0.31* | 0.66 | 0.53 | MACR not resolved from overlying 70 |
| 9/28/97 | 0609-0900 | 0.22 | | 0.31 | | 0.39 | | |
| 9/28/97 | 0904-1200 | 0.50 | 0.50 | 0.42 | 0.37 | 0.99 | 0.83 | |
| 9/28/97 | 0904-1200 | 0.50 | | 0.32 | | 0.66 | | |
| 9/28/97 | 1300-1600 | 0.44 | 0.42 | 0.41 | 0.41 | 0.66 | 0.68 | |
| 9/28/97 | 1300-1600 | 0.39 | | 0.40 | | 0.69 | | |
| 9/28/97 | 1700-2000 | 0.60 | 0.34* | 0.78 | 0.25* | 1.23 | 0.31* | Very low isoprene-d ₈ |
| 9/28/97 | 1700-2000 | 0.34 | | 0.25 | | 0.31 | | |
| 9/28-29/97 | 2015-0300 | 0.06 | 0.07 | 0.19 | 0.19 | 0.16 | 0.23 | |
| 9/28-29/97 | 2015-0300 | 0.08 | | 0.18 | | 0.30 | | |
| 9/29/97 | 0310-0600 | 0.13 | 0.13 | 0.38 | 0.36 | 0.34 | 0.29 | |
| 9/29/97 | 0310-0600 | 0.12 | | 0.33 | | 0.23 | | |
| 9/29/97 | 0616-0858 | 0.31 | 0.30 | 0.45 | 0.44 | 0.45 | 0.65 | |
| 9/29/97 | 0616-0858 | 0.30 | | 0.43 | | 0.86 | | |
| 9/29/97 | 0900-1200 | 0.38 | 0.37 | 0.37 | 0.30 | 0.60 | 0.46 | |
| 9/29/97 | 0900-1200 | 0.36 | | 0.24 | | 0.32 | | |
| 9/29/97 | 1300-1600 | 0.38 | 0.38 | 0.31 | 0.61 | 0.62 | 0.74 | |
| 9/29/97 | 1300-1600 | 0.37 | | 0.92 | | 0.85 | | |
| 9/29/97 | 1700-2000 | 0.40 | 0.44 | 0.22 | 0.22 | 0.44 | 0.40 | |
| 9/29/97 | 1700-2000 | 0.48 | | 0.21 | | 0.36 | | |

Table A-1.1. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Azusa
(continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|-----------|---------------|------------------|----------------------|--------------|------------------|-------------|-----------------|--|
| AZUSA | Oct 3-4, 1997 | | | | | | | |
| 10/3/97 | 0301-0600 | 0.05 | 0.06 | 0.12 | 0.13 | 0.31 | 0.26 | |
| 10/3/97 | 0301-0600 | 0.07 | | 0.14 | | 0.21 | | |
| 10/3/97 | 0614-0900 | 0.06 | 0.06 | 0.08 | 0.07 | 0.11 | 0.12 | |
| 10/3/97 | 0614-0900 | 0.06 | | 0.06 | | 0.12 | | |
| 10/3/97 | 0904-1200 | 0.12 | 0.12 | 0.10 | 0.10 | 0.21 | 0.22 | |
| 10/3/97 | 0904-1200 | 0.12 | | 0.09 | | 0.24 | | |
| 10/3/97 | 1300-1600 | 0.21 | 0.21 | 0.21 | 0.19 | 0.42 | 0.38 | |
| 10/3/97 | 1300-1600 | 0.21 | | 0.16 | | 0.34 | | |
| 10/3/97 | 1700-2000 | 0.17 | 0.15 | 0.18 | 0.15 | 0.53 | 0.44 | |
| 10/3/97 | 1700-2000 | 0.13 | | 0.12 | | 0.35 | | |
| 10/3-4/97 | 2011-0300 | 0.06 | 0.06 | 0.21 | 0.19 | 0.66 | 0.50 | |
| 10/3-4/97 | 2011-0300 | 0.05 | | 0.18 | | 0.33 | | |
| 10/4/97 | 0304-0600 | 0.12 | 0.12 | 0.18 | 0.18 | 0.98 | 0.99 | |
| 10/4/97 | 0304-0600 | 0.13 | | 0.19 | | 1.01 | | |
| 10/5/97 | 0612-0900 | 0.07 | 0.07 | 0.07 | 0.08 | 0.24 | 0.26 | |
| 10/4/97 | 0612-0900 | 0.07 | | 0.10 | | 0.29 | | |
| 10/4/97 | 0901-1200 | 0.17 | 0.16 | 0.17 | 0.16 | 0.35 | 0.36 | |
| 10/4/97 | 0901-1200 | 0.16 | | 0.16 | | 0.38 | | |
| 10/4/97 | 1300-1600 | 0.36 | 0.24* | 0.51 | 0.29* | 1.06 | 0.64* | Sampling tube contents transferred from broken tube |
| 10/4/97 | 1300-1600 | 0.24 | | 0.29 | | 0.64 | | |
| 10/4/97 | 1700-2000 | 0.14 | 0.14 | 0.12 | 0.11 | 0.22 | 0.22 | |
| 10/4/97 | 1700-2000 | 0.14 | | 0.11 | | 0.23 | | |

*Used single value.

Table A-1.2. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Pine Mtn. and Mt. Baldy.

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|------------------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|----------------------------|
| PINE MTN. | | | | | | | | |
| Aug 4-6, 1997 | | | | | | | | |
| 8/4/97 | 1710-2000 | 2.12 | 2.21 | 0.89 | 0.91 | 1.54 | 1.58 | |
| 8/4/97 | 1710-2000 | 2.30 | | 0.94 | | 1.63 | | |
| 08/4-08/5/97 | 2025-0259 | 0.04 | 0.03 | 0.60 | 0.58 | 1.15 | 1.11 | |
| 08/4-08/5/97 | 2025-0259 | 0.02 | | 0.55 | | 1.07 | | |
| 8/5/97 | 0314-0559 | 0.05 | 0.04 | 0.15 | 0.14 | 0.26 | 0.25 | |
| 8/5/97 | 0314-0559 | 0.03 | | 0.13 | | 0.24 | | |
| 8/5/97 | 0617-0858 | 1.85 | 1.96 | 0.09 | 0.09 | 0.15 | 0.16 | |
| 8/5/97 | 0617-0858 | 2.08 | | 0.09 | | 0.17 | | |
| 8/5/97 | 0911-1200 | 0.96 | 1.00 | 0.30 | 0.33 | 0.81 | 0.83 | |
| 8/5/97 | 0911-1200 | 1.05 | | 0.35 | | 0.85 | | |
| 8/5/97 | 1302-1600 | 0.59 | 0.59* | 0.46 | 0.46* | 0.71 | 0.71* | |
| 8/5/97 | 1302-1600 | - | | - | | - | | Sampling tube missing |
| 8/5/97 | 1700-2000 | 1.04 | 1.09 | 0.42 | 0.44 | 0.68 | 0.72 | |
| 8/5/97 | 1700-2000 | 1.15 | | 0.47 | | 0.77 | | |
| 8/5/97 | 2015-0255 | 0.08 | 0.08 | 0.07 | 0.08 | 0.18 | 0.19 | |
| 8/5/97 | 2015-0255 | 0.08 | | 0.08 | | 0.20 | | |
| 8/6/97 | 0310-0600 | 0.03 | 0.03 | 0.03 | 0.03 | 0.07 | 0.07 | |
| 8/6/97 | 0310-0600 | 0.04 | | 0.03 | | 0.07 | | |
| 8/6/97 | 0615-0900 | 0.76 | 0.80 | 0.07 | 0.07 | 0.14 | 0.14 | |
| 8/6/97 | 0615-0900 | 0.84 | | 0.07 | | 0.15 | | |
| 8/6/97 | 0915-1200 | 0.41 | 0.41 | 0.18 | 0.19 | 0.38 | 0.36 | |
| 8/6/97 | 0915-1200 | 0.41 | | 0.20 | | 0.34 | | |
| 8/6/97 | 1300-1600 | 0.58 | 0.62 | 0.54 | 0.59 | 0.92 | 1.03 | |
| 8/6/97 | 1300-1600 | 0.65 | | 0.65 | | 1.13 | | |

Table A-1.2. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Pine Mtn. and Mt. Baldy (continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|------------------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|----------------------------|
| PINE MTN. | | | | | | | | |
| Sep 4-7, 1997 | | | | | | | | |
| 9/4/97 | 1702-2000 | 0.52 | 0.41 | 0.72 | 0.52 | 0.86 | 0.72 | |
| 9/4/97 | 1702-2000 | 0.31 | | 0.32 | | 0.57 | | |
| 9/4/97 | 2008-2400 | 0.03 | 0.03 | 0.35 | 0.33 | 0.57 | 0.54 | |
| 9/4/97 | 2008-2400 | 0.03 | | 0.32 | | 0.52 | | |
| 9/5/97 | 0010-0338 | 0.03 | 0.02 | 0.10 | 0.10 | 0.23 | 0.16 | |
| 9/5/97 | 0010-0338 | 0.02 | | 0.10 | | 0.09 | | |
| 9/5/97 | 0349-0600 | 0.01 | 0.02 | 0.08 | 0.09 | 0.17 | 0.18 | |
| 9/5/97 | 0349-0600 | 0.02 | | 0.09 | | 0.20 | | |
| 9/5/97 | 0615-0900 | 0.66 | 0.63 | 0.10 | 0.10 | 0.18 | 0.20 | |
| 9/5/97 | 0615-0900 | 0.60 | | 0.11 | | 0.21 | | |
| 9/5/97 | 0908-1200 | 0.96 | 0.93 | 0.21 | 0.21 | 0.38 | 0.38 | |
| 9/5/97 | 0908-1200 | 0.89 | | 0.21 | | 0.38 | | |
| 9/5/97 | 1300-1600 | 1.73 | 1.71 | 1.05 | 1.07 | 0.96 | 1.38 | |
| 9/5/97 | 1300-1600 | 1.68 | | 1.09 | | 1.79 | | |
| 9/5/97 | 1700-2000 | 0.62 | 0.59 | 0.31 | 0.30 | 0.53 | 0.49 | |
| 9/5/97 | 1700-2000 | 0.55 | | 0.28 | | 0.46 | | |
| 9/5/97 | 2007-2355 | 0.04 | 0.03 | 0.21 | 0.26 | 0.32 | 0.39 | |
| 9/5/97 | 2007-2355 | 0.02 | | 0.31 | | 0.46 | | |
| 9/6/97 | 0000-0255 | 0.02 | 0.02 | 0.09 | 0.10 | 0.18 | 0.17 | |
| 9/6/97 | 0000-0255 | 0.02 | | 0.10 | | 0.16 | | |
| 9/6/97 | 0301-0600 | 0.03 | 0.03 | 0.11 | 0.10 | 0.17 | 0.14 | |
| 9/6/97 | 0301-0600 | 0.03 | | 0.08 | | 0.11 | | |
| 9/6/97 | 0610-0900 | 0.52 | 0.48 | 0.08 | 0.08 | 0.17 | 0.16 | |
| 9/6/97 | 0610-0900 | 0.44 | | 0.08 | | 0.15 | | |
| 9/6/97 | 0905-1200 | 0.56 | 0.60 | 0.20 | 0.17 | 0.04 | 0.17 | |
| 9/6/97 | 0905-1200 | 0.64 | | 0.15 | | 0.30 | | |

Table A-1.2. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Pine Mtn. and Mt. Baldy (continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|------------------|------------------------------|------------------|----------------------|--------------|------------------|-------------|-----------------|----------------------------------|
| 9/6/97 | 1300-1600 | 1.62 | 1.58 | 0.36 | 0.25 | 0.45 | 0.35 | |
| 9/6/97 | 1300-1600 | 1.54 | | 0.15 | | 0.25 | | |
| 9/6/97 | 1700-2000 | 0.56 | 0.55 | 0.33 | 0.33 | 0.54 | 0.54 | |
| 9/6/97 | 1700-2000 | 0.54 | | 0.34 | | 0.54 | | |
| 9/6/97 | 2007-2355 | 0.06 | 0.05 | 0.17 | 0.18 | 0.28 | 0.26 | |
| 9/6/97 | 2007-2355 | 0.03 | | 0.18 | | 0.23 | | |
| 9/7/97 | 0000-0255 | 0.03 | 0.03 | 0.18 | 0.16 | 0.23 | 0.24 | |
| 9/7/97 | 0000-0255 | 0.03 | | 0.15 | | 0.25 | | |
| 9/7/97 | 0300-0555 | 0.05 | 0.05 | 0.13 | 0.10 | 0.19 | 0.15 | |
| 9/7/97 | 0300-0555 | 0.04 | | 0.08 | | 0.11 | | |
| 9/7/97 | 0600-0900 | 1.83 | 1.88 | 0.12 | 0.10 | 0.27 | 0.21 | |
| 9/7/97 | 0600-0900 | 1.92 | | 0.07 | | 0.15 | | |
| MT. BALDY | | | | | | | | |
| 9/28/97 | SEP 28-29, 1997 0904-1200 | 1.91 | 1.90 | 0.70 | 0.23* | 0.32 | 0.40 | (Ion 69 used to quantitate MACR) |
| 9/28/97 | 0904-1200 | 1.90 | | 0.23 | | 0.48 | | (Ion 69 used to quantitate MACR) |
| 9/28/97 | 1300-1600 | 1.14 | 1.17 | 0.37 | 0.37 | 0.91 | 0.93 | |
| 9/28/97 | 1300-1600 | 1.21 | | 0.36 | | 0.95 | | |
| 9/28/97 | 1700-2000 | 2.15 | 2.27 | 0.39 | 0.40 | 0.82 | 0.85 | |
| 9/28/97 | 1700-2000 | 2.38 | | 0.40 | | 0.88 | | |
| 9/28-29/97 | 2005-0300 | 0.09 | 0.08 | 0.24 | 0.24 | 0.44 | 0.47 | |
| 9/28-29/97 | 2005-0300 | 0.07 | | 0.23 | | 0.50 | | |
| 9/29/97 | 0307-0600 | 0.06 | 0.06 | 0.07 | 0.06 | 0.17 | 0.17 | |
| 9/29/97 | 0307-0600 | 0.06 | | 0.06 | | 0.17 | | |
| 9/29/97 | 0606-0900 | 0.69 | 0.65 | 0.06 | 0.06 | 0.15 | 0.15 | |
| 9/29/97 | 0606-0900 | 0.62 | | 0.06 | | 0.15 | | |
| 9/29/97 | 0903-1200 | 1.84 | 1.84 | 0.28 | 0.27 | 0.65 | 0.63 | |
| 9/29/97 | 0903-1200 | 1.85 | | 0.26 | | 0.61 | | |

Table A-1.2. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Pine Mtn. and Mt. Baldy (continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|--------------------------------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|----------------------------|
| 9/29/97 | 1300-1600 | 1.08 | 1.07 | 0.29 | 0.29 | 0.60 | 0.61 | |
| 9/29/97 | 1300-1600 | 1.07 | | 0.28 | | 0.61 | | |
| 9/29/97 | 1700-2000 | 1.94 | 1.92 | 0.44 | 0.39 | 0.91 | 0.79 | |
| 9/29/97 | 1700-2000 | 1.89 | | 0.34 | | 0.66 | | |
| MT. BALDY OCT 3-4, 1997 | | | | | | | | |
| 10/3/97 | 0619-0900 | 0.42 | 0.42 | 0.02 | 0.02 | 0.11 | 0.12 | |
| 10/3/97 | 0619-0900 | 0.41 | | 0.02 | | 0.13 | | |
| 10/3/97 | 0903-1200 | 1.39 | 1.38 | 0.13 | 0.13 | 0.28 | 0.28 | |
| 10/3/97 | 0903-1200 | 1.37 | | 0.12 | | 0.28 | | |
| 10/3/97 | 1301-1600 | 1.22 | 1.23 | 0.41 | 0.48 | 0.77 | 1.01 | |
| 10/3/97 | 1301-1600 | 1.23 | | 0.55 | | 1.25 | | |
| 10/3/97 | 1703-2003 | 1.29 | 1.29* | 0.37 | 0.37* | 0.72 | 0.72* | |
| 10/3/97 | 1703-2003 | 1.91 | | 0.60 | | 1.07 | | |
| 10/3-4/97 | 2010-0302 | 0.05 | 0.05 | 0.12 | 0.13 | 0.24 | 0.26 | |
| 10/3-4/97 | 2010-0302 | 0.05 | | 0.13 | | 0.27 | | |
| 10/4/97 | 0310-0601 | 0.04 | 0.04 | 0.06 | 0.06 | 0.11 | 0.11 | |
| 10/4/97 | 0310-0601 | 0.04 | | 0.05 | | 0.12 | | |
| 10/4/97 | 0612-0900 | 0.26 | 0.26 | 0.04 | 0.04 | 0.10 | 0.10 | |
| 10/4/97 | 0612-0900 | 0.27 | | 0.04 | | 0.10 | | |
| 10/4/97 | 0903-1200 | 1.54 | 1.53 | 0.11 | 0.12 | 0.24 | 0.25 | |
| 10/4/97 | 0903-1200 | 1.51 | | 0.12 | | 0.25 | | |
| 10/4/97 | 1300-1600 | 0.86 | 0.88 | 0.33 | 0.32 | 0.77 | 0.75 | |
| 10/4/97 | 1300-1600 | 0.90 | | 0.31 | | 0.73 | | |
| 10/4/97 | 1700-2000 | 1.30 | 1.29 | 0.36 | 0.36 | 0.67 | 0.66 | |

*Used single value.

Table A-1.3. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Banning

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|------------------------------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|---|
| BANNING AUG 4-6, 1997 | | | | | | | | |
| 8/4/97 | 0600-0855 | 0.21 | 0.25 | 0.12 | 0.14 | 0.25 | 0.33 | |
| 8/4/97 | 0600-0855 | 0.29 | | 0.16 | | 0.40 | | |
| 8/4/97 | 0902-1157 | 0.06 | 0.06 | 0.05 | 0.09 | 0.16 | 0.19 | |
| 8/4/97 | 0902-1157 | 0.07 | | 0.12 | | 0.23 | | |
| 8/4/97 | - | - | - | - | - | - | - | Wind knocked over sampling equipment, no sample |
| 8/4/97 | - | - | - | - | - | - | - | Wind knocked over sampling equipment, no sample |
| 8/4/97 | 1504-1800 | 0.18 | 0.21 | 0.15 | 0.12 | 0.34 | 0.28 | |
| 8/4/97 | 1504-1800 | 0.24 | | 0.09 | | 0.21 | | |
| 8/4-5/1997 | 1807-0255 | 0.08 | 0.08 | 0.40 | | 0.35 | | |
| 8/4-5/1997 | 1807-0255 | 0.08 | | 0.30 | | 0.59 | | |
| 8/5/97 | 0303-0550 | 0.02 | 0.02 | 0.16 | 0.16 | 0.25 | 0.26 | |
| 8/5/97 | 0303-0550 | 0.02 | | 0.16 | | 0.27 | | |
| 8/5/97 | 0604-0855 | 0.32 | 0.33 | 0.21 | 0.20 | 0.30 | 0.31 | |
| 8/5/97 | 0604-0855 | 0.34 | | 0.20 | | 0.32 | | |
| 8/5/97 | 0901-1153 | 0.07 | 0.08 | 0.07 | 0.07 | 0.15 | 0.15 | |
| 8/5/97 | 0901-1153 | 0.08 | | 0.06 | | 0.15 | | |
| 8/5/97 | 1159-1452 | 0.08 | 0.10 | 0.04 | 0.04 | 0.10 | 0.11 | |
| 8/5/97 | 1159-1452 | 0.12 | | 0.04 | | 0.12 | | |
| 8/5/97 | 1458-1800 | 0.12 | 0.13 | 0.07 | 0.07 | 0.12 | 0.13 | |
| 8/5/97 | 1458-1800 | 0.14 | | 0.07 | | 0.14 | | |
| 8/5- 8/6/97 | 1816-0255 | 0.06 | 0.07 | 0.15 | 0.22 | 0.30 | 0.39 | |
| 8/5- 8/6/97 | 1816-0255 | 0.08 | | 0.28 | | 0.48 | | |
| 8/6/97 | 0301-0550 | 0.01 | 0.01 | 0.03 | 0.04 | 0.07 | 0.10 | |
| 8/6/97 | 0301-0550 | 0.02 | | 0.06 | | 0.14 | | |
| 8/6/97 | 0605-0855 | 0.02 | | 0.02 | 0.02 | 0.04 | 0.04 | |
| 8/6/97 | 0605-0855 | 0.02 | | 0.02 | | 0.04 | | |

Table A-1.3. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Banning
(continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | Isoprene ppbv | MACR av, ppbv | MACR av, ppbv | MVK av, ppbv | MVK av, ppbv | Reason for discarding data |
|--------------------------------|-----------|------------------|----------------------|------------------|------------------|------------------|-----------------|-----------------|----------------------------|
| BANNING AUG 22-23, 1997 | | | | | | | | | |
| 8/22/97 | 0901-1150 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.04 | 0.04 | 0.05 |
| 8/6/97 | 0901-1150 | 0.03 | 0.03 | 0.03 | 0.02 | 0.02 | 0.05 | 0.05 | 0.06 |
| 8/6/97 | 1155-1455 | 0.04 | 0.04 | 0.03 | 0.02 | 0.02 | 0.05 | 0.05 | 0.07 |
| 8/6/97 | 1155-1455 | 0.03 | 0.03 | 0.02 | 0.02 | 0.02 | 0.07 | 0.07 | 0.14 |
| 8/6/97 | 1500-1800 | 0.16 | 0.18 | 0.04 | 0.05 | 0.05 | 0.14 | 0.16 | 0.17 |
| 8/6/97 | 1500-1800 | 0.19 | 0.05 | | | | | | |
| BANNING AUG 22-23, 1997 | | | | | | | | | |
| 8/22/97 | 0600-0855 | 0.11 | 0.12 | 0.06 | 0.07 | 0.07 | 0.31 | 0.31 | 0.36 |
| 8/22/97 | 0600-0855 | 0.12 | 0.08 | 0.08 | 0.10 | 0.10 | 0.41 | 0.41 | 0.26 |
| 8/22/97 | 0900-1155 | 0.24 | 0.27 | 0.10 | 0.10 | 0.10 | 0.25 | 0.27 | |
| 8/22/97 | 0900-1155 | 0.29 | 0.20 | 0.09 | 0.09 | 0.09 | 0.22 | 0.22 | 0.23 |
| 8/22/97 | 1200-1455 | 0.19 | 0.20 | 0.09 | 0.09 | 0.09 | 0.23 | 0.23 | 0.26 |
| 8/22/97 | 1200-1455 | 0.21 | 0.28 | 0.12 | 0.12 | 0.14 | 0.32 | 0.32 | |
| 8/22/97 | 1500-1800 | 0.28 | 0.28 | 0.17 | 0.17 | 0.17 | 0.37 | 0.37 | |
| 8/22-23/97 | 1500-1800 | 0.28 | 0.08 | 2.45 | 2.45 | 2.37 | 3.08 | 3.08 | 2.93 |
| 8/22-23/97 | 1813-0300 | 0.09 | 0.07 | 2.29 | 2.29 | 2.29 | 2.79 | 2.79 | |
| 8/22-23/97 | 1813-0300 | 0.07 | 0.01 | 0.13 | 0.13 | 0.16 | 0.32 | 0.32 | |
| 8/23/97 | 0303-0555 | 0.01 | 0.14 | 0.18 | 0.18 | 0.12 | 0.51 | 0.51 | |
| 8/23/97 | 0303-0555 | 0.02 | 0.14 | 0.14 | 0.12 | 0.12 | 0.26 | 0.26 | 0.27 |
| 8/23/97 | 0603-0900 | 0.14 | 0.14 | 0.12 | 0.12 | 0.12 | 0.27 | 0.27 | |
| 8/23/97 | 0603-0900 | 0.14 | 0.01 | 0.16 | 0.16 | 0.07 | 0.16 | 0.16 | 0.42 |
| 8/23/97 | 0903-1200 | 0.20 | 0.22 | 0.06 | 0.06 | 0.07 | 0.20 | 0.20 | |
| 8/23/97 | 0903-1200 | 0.24 | 0.20 | 0.08 | 0.08 | 0.05 | 0.12 | 0.12 | 0.12 |
| 8/23/97 | 1202-1500 | 0.19 | 0.20 | 0.05 | 0.05 | 0.05 | 0.12 | 0.12 | |
| 8/23/97 | 1202-1500 | 0.21 | 0.05 | 0.05 | 0.05 | 0.05 | 0.12 | 0.12 | |
| 8/23/97 | 1502-1800 | 0.20 | 0.22 | 0.08 | 0.08 | 0.07 | 0.25 | 0.25 | |
| 8/23/97 | 1502-1800 | 0.23 | 0.07 | | | | 0.25 | 0.25 | |

Table A-1.3. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Banning
(continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|------------------------------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|----------------------------|
| BANNING SEP 5-6, 1997 | | | | | | | | |
| 9/5/97 | 0600-0855 | 0.12 | 0.12 | 0.20 | 0.21 | 0.44 | 0.39 | |
| 9/5/97 | 0600-0855 | 0.13 | | 0.21 | | 0.34 | | |
| 9/5/97 | 0900-1155 | 0.18 | 0.19 | 0.09 | 0.09 | 0.19 | 0.21 | |
| 9/5/97 | 0900-1155 | 0.19 | | 0.09 | | 0.22 | | |
| 9/5/97 | 1158-1458 | 0.16 | 0.17 | 0.04 | 0.03 | 0.11 | 0.11 | |
| 9/5/97 | 1158-1458 | 0.18 | | 0.03 | | 0.10 | | |
| 9/5/97 | 1501-1800 | 0.17 | 0.17 | 0.18 | 0.18 | 0.59 | 0.59 | |
| 9/5/97 | 1501-1800 | 0.18 | | 0.18 | | 0.60 | | |
| 9/5-6/97 | 1824-0257 | 0.03 | 0.05 | 0.35 | 0.45 | 0.50 | 0.53 | |
| 9/5-6/97 | 1824-0257 | 0.06 | | 0.55 | | 0.57 | | |
| 9/6/97 | 0308-0550 | 0.01 | 0.01* | 0.29 | 0.29* | 0.29 | 0.29* | |
| 9/6/97 | 0308-0550 | - | - | - | - | - | - | |
| 9/6/97 | 0610-0855 | 0.01 | 0.10* | 0.11 | 0.33* | 0.19 | 0.44* | |
| 9/6/97 | 0610-0855 | 0.10 | | 0.33 | | 0.44 | | |
| 9/6/97 | 0900-1158 | 0.03 | | 0.08 | 0.05 | 0.16 | 0.14 | |
| 9/6/97 | 0900-1158 | 0.02 | | 0.03 | | 0.13 | | |
| 9/6/97 | 1202-1458 | 0.11 | | 0.12 | 0.04 | 0.04 | 0.11 | |
| 9/6/97 | 1205-1458 | 0.13 | | 0.04 | | 0.10 | | |
| 9/6/97 | 1502-1800 | 0.21 | 0.21* | 0.06 | 0.06* | 0.18 | 0.18* | |
| 9/6/97 | 1502-1800 | 1.64 | | 0.60 | | 1.85 | | |
| BANNING OCT 3-4, 1997 | | | | | | | | |
| 10/3/97 | 0600-0858 | 0.05 | 0.05 | 0.07 | 0.08 | 0.27 | 0.25 | |
| 10/3/97 | 0600-0858 | 0.05 | | 0.10 | | 0.22 | | |
| 10/3/97 | 0900-1158 | 0.06 | | 0.05 | 0.07 | 0.14 | 0.14 | |
| 10/3/97 | 0900-1158 | 0.06 | | 0.08 | | 0.14 | | |

Sampling tube contents expelled from tube
desorber

Results low, seems like a back sampling tube

Table A-1.3. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Bamming (continued).

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|-----------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|----------------------------|
| 10/3/97 | 1200-1458 | 0.03 | 0.03 | 0.03 | 0.03 | 0.10 | 0.08 | |
| 10/3/97 | 1200-1458 | 0.03 | 0.03 | 0.03 | 0.03 | 0.07 | | |
| 10/3/97 | 1500-1800 | 0.19 | 0.19 | 0.15 | 0.15 | 0.36 | 0.36 | |
| 10/3/97 | 1500-1800 | 0.20 | | 0.15 | | 0.35 | | |
| 10/3-4/97 | 1807-0258 | 0.04 | 0.04 | 0.17 | 0.14 | 0.17 | 0.18 | |
| 10/3-4/97 | 1807-0258 | 0.03 | | 0.11 | | 0.18 | | |
| 10/4/97 | 0300-0553 | 0.03 | 0.04 | 0.19 | 0.18 | 0.30 | 0.32 | |
| 10/4/97 | 0300-0553 | 0.04 | | 0.18 | | 0.33 | | |
| 10/4/97 | 0601-0858 | 0.10 | 0.10 | 0.17 | 0.16 | 0.32 | 0.29 | |
| 10/4/97 | 0601-0858 | 0.10 | | 0.16 | | 0.27 | | |
| 10/4/97 | 0900-1158 | 0.14 | 0.14 | 0.10 | 0.10 | 0.32 | 0.30 | |
| 10/4/97 | 0900-1158 | 0.13 | | 0.10 | | 0.27 | | |
| 10/4/97 | 1200-1458 | 0.13 | 0.12 | 0.11 | 0.10 | 0.31 | 0.28 | |
| 10/4/97 | 1200-1458 | 0.12 | | 0.09 | | 0.24 | | |
| 10/4/97 | 1500-1800 | 0.10 | 0.11 | 0.13 | 0.16 | 0.45 | 0.46 | |
| 10/4/97 | 1500-1800 | 0.13 | | 0.19 | | 0.47 | | |

*Used single value.

Table A-1.4. Detailed Data from Measurements of Isoprene and its Atmospheric Reaction Products in Ambient Air at Los Angeles,
North Main Street.

| Date | Sampling | Isoprene ppbv | Isoprene av, ppbv | MACR ppbv | MACR av, ppbv | MVK ppbv | MVK av, ppbv | Reason for discarding data |
|------------------------------------|-----------|------------------|----------------------|--------------|------------------|-------------|-----------------|----------------------------|
| LOS ANGELES, NORTH MAIN ST. | | | | | | | | |
| 8/22/97 | 0558-0855 | 0.40 | 0.37 | 0.75 | 0.77 | 0.53 | 0.50 | |
| 8/22/97 | 0558-0855 | 0.35 | | 0.80 | | 0.47 | | |
| 8/22/97 | 0900-1155 | 0.10 | 0.11 | 1.12 | 1.26 | 1.10 | 0.98 | |
| 8/22/97 | 0900-1155 | 0.11 | | 1.41 | | 0.85 | | |
| 8/22/97 | 1202-1457 | 0.22 | 0.21 | 0.50 | 0.49 | 0.72 | 0.67 | |
| 8/22/97 | 1202-1457 | 0.19 | | 0.48 | | 0.62 | | |
| 8/22/97 | 1500-1755 | 0.14 | 0.17 | 0.22 | 0.27 | 0.41 | 0.47 | |
| 8/22/97 | 1500-1755 | 0.19 | | 0.33 | | 0.54 | | |
| 8/22-23/97 | 1806-0556 | 0.19 | 0.23 | 0.22 | 0.28 | 0.20 | 0.41 | |
| 8/22-23/97 | 1806-0556 | 0.28 | | 0.35 | | 0.61 | | |
| 8/23/97 | 0605-0857 | 0.18 | 0.21 | 0.34 | 0.44 | 0.32 | 0.44 | |
| 8/23/97 | 0605-0857 | 0.24 | | 0.54 | | 0.56 | | |
| 8/23/97 | 0900-1156 | 0.13 | 0.12 | 0.34 | 0.33 | 0.62 | 0.73 | |
| 8/23/97 | 0900-1156 | 0.12 | | 0.32 | | 0.83 | | |
| 8/23/97 | 1202-1458 | 0.14 | 0.15 | 0.36 | 0.39 | 0.85 | 0.82 | |
| 8/23/97 | 1202-1458 | 0.16 | | 0.42 | | 0.79 | | |
| 8/23/97 | 1500-1800 | 0.21 | 0.21 | 0.30 | 0.27 | 0.45 | 0.50 | |
| 8/23/97 | 1500-1800 | 0.21 | | 0.24 | | 0.54 | | |

Table B-1.1. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Azusa.

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|----------------------------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|----------------------------|
| AZUSA AUG 4-6, 1997 | | | | | | | | | | |
| 8/4/97 | 0600-0900 | 0.3122 | 0.3196 | 0.0100 | 0.0108 | 0.3304 | 0.3454 | 0.0432 | 0.0451 | |
| 8/4/97 | 0600-0900 | 0.3271 | 0.0115 | 0.0052 | 0.0054 | 0.3604 | 0.0791 | 0.0470 | 0.0470 | |
| 8/4/97 | 0907-1206 | 0.1061 | 0.1106 | 0.0057 | 0.0057 | 0.1267 | 0.0104 | 0.0156 | 0.0162 | |
| 8/4/97 | 0907-1206 | 0.1150 | 0.0284 | 0.0278 | 0.0039 | 0.0029 | 0.0155 | 0.0169 | 0.0075 | 0.0080 |
| 8/4/97 | 1306-1603 | 0.0272 | 0.019 | 0.019 | 0.0019 | 0.0206 | 0.0206 | 0.0084 | 0.0084 | |
| 8/4/97 | 1306-1603 | 0.0469 | 0.0446 | 0.0030 | 0.0043 | 0.0416 | 0.0283 | 0.0161 | 0.0148 | |
| 8/4/97 | 1700-2000 | 0.0423 | 0.0056 | 0.0085 | 0.0080 | 0.0149 | 0.0149 | 0.0135 | 0.0135 | |
| 8/4/97 | 2006-0300 | 0.1778 | 0.1800 | 0.0074 | 0.0074 | 0.1569 | 0.1526 | 0.0369 | 0.0368 | |
| 8/4/97 | 2006-0300 | 0.1822 | 0.3515 | 0.3768 | 0.0196 | 0.1482 | 0.1482 | 0.0366 | 0.0366 | |
| 8/5/97 | 0307-0600 | 0.4020 | 0.0225 | 0.2187 | 0.2214 | 0.0133 | 0.0137 | 0.0925 | 0.0925 | |
| 8/5/97 | 0307-0600 | 0.2187 | 0.2241 | 0.0141 | 0.0141 | 0.0877 | 0.0877 | 0.0588 | 0.0588 | |
| 8/5/97 | 0618-0900 | 0.1204 | 0.1221 | 0.0066 | 0.0064 | 0.1041 | 0.1041 | 0.0644 | 0.0644 | |
| 8/5/97 | 0907-1202 | 0.1238 | 0.0063 | 0.0336 | 0.0332 | 0.0037 | 0.0041 | 0.1191 | 0.1191 | |
| 8/5/97 | 0907-1202 | 0.1238 | 0.0336 | 0.0332 | 0.0332 | 0.0183 | 0.0183 | 0.0109 | 0.0109 | |
| 8/5/97 | 1300-1600 | 0.0328 | 0.045 | 0.0465 | 0.0470 | 0.0042 | 0.0037 | 0.0264 | 0.0283 | 0.0198 |
| 8/5/97 | 1300-1600 | 0.0328 | 0.0465 | 0.0475 | 0.0475 | 0.0031 | 0.0031 | 0.0301 | 0.0204 | |
| 8/5/97 | 1701-2000 | 0.0465 | 0.0470 | 0.2825 | 0.2977 | 0.0128 | 0.0134 | 0.2358 | 0.2617 | 0.0109 |
| 8/5- 8/6/97 | 2005-0300 | 0.3130 | 0.4373 | 0.4534 | 0.0152 | 0.0164 | 0.0287 | 0.0283 | 0.0160 | 0.0158 |
| 8/6/97 | 0306-0600 | 0.4695 | 0.3490 | 0.3490 | 0.0176 | 0.0174 | 0.0170 | 0.0301 | 0.0156 | 0.0109 |
| 8/6/97 | 0616-0900 | 0.3715 | 0.0167 | 0.0167 | 0.0167 | 0.2539 | 0.2539 | 0.2061 | 0.0478 | 0.0499 |
| 8/6/97 | 0616-0900 | 0.3264 | 0.0789 | 0.0808 | 0.0076 | 0.0068 | 0.0308 | 0.0371 | 0.0581 | 0.0581 |
| 8/6/97 | 0906-1200 | 0.0827 | 0.0059 | 0.0059 | 0.0059 | 0.0434 | 0.0434 | 0.0158 | 0.0161 | 0.0163 |
| 8/6/97 | 0906-1200 | 0.0827 | 0.0059 | 0.0059 | 0.0059 | 0.0163 | 0.0163 | 0.0163 | 0.0163 | |

Table B-1.1. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Azusa (continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|--------------|------------------------------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|---|
| 8/6/97 | 1300-1600 | 0.0390 | 0.0390 | 0.0029 | 0.0028 | 0.0159 | 0.0159 | 0.0085 | 0.0087 | |
| 8/6/97 | 1300-1600 | 0.0390 | 0.0026 | | | 0.0158 | | 0.0090 | | |
| 8/6/97 | 1700-2000 | 0.0314 | 0.0322 | 0.0022 | 0.0021 | 0.0186 | 0.0147 | 0.0102 | 0.0096 | |
| 8/6/97 | 1700-2000 | 0.0330 | | 0.0021 | | 0.0109 | | 0.0091 | | |
| AZUSA | | | | | | | | | | |
| 8/22/97 | Aug 22-23, 1997 0318-0600 | 0.1675* | 0.1675* | 0.1396 | 0.1396* | 0.0178 | 0.0178* | 0.0073 | 0.0073* | |
| 8/22/97 | 0318-0600 | 0.8046 | | 0.3350 | | 0.0499 | | 0.0419 | | |
| 8/22/97 | 0610-0900 | 0.0010 | 0.3739* | 0.0013 | 0.0084* | 0.0013 | 0.2562* | ** | 0.0602* | |
| 8/22/97 | 0610-0900 | 0.3739 | | 0.0084 | | 0.2562 | | 0.0602 | | |
| 8/22/97 | 0905-1157 | 0.1204 | 0.1051 | 0.0067 | 0.0128 | 0.0243 | 0.0155 | 0.0227 | 0.0173 | |
| 8/22/97 | 0905-1157 | 0.0897 | | 0.0189 | | 0.0067 | | 0.0120 | | |
| 8/22/97 | 1210-1500 | 0.0261 | 0.0395* | 0.0223 | 0.0074* | ** | 0.0024* | 0.0028 | 0.0035* | |
| 8/22/97 | 1210-1500 | 0.0395 | | 0.0074 | | 0.0024 | | 0.0035 | | |
| 8/22/97 | 1504-1800 | 0.0327 | 0.0340 | 0.0042 | 0.0052 | 0.0084 | 0.0050 | 0.0101 | 0.0078 | |
| 8/22/97 | 1504-1800 | 0.0352 | | 0.0061 | | 0.0016 | | 0.0054 | | |
| 8/22-23/97 | 1815-0300 | 0.1490 | 0.1429 | 0.0063 | 0.0086 | 0.0713 | 0.0496 | 0.0279 | 0.0246 | |
| 8/22-23/97 | 1815-0300 | 0.1369 | | 0.0110 | | 0.0279 | | 0.0213 | | |
| 8/23/97 | 0304-0600 | 0.2552 | 0.4371* | 0.0882 | 0.0113* | 0.0180 | 0.2606* | 0.0169 | 0.0666* | Altered monoterpene profile |
| 8/23/97 | 0304-0600 | 0.4371 | | 0.0113 | | 0.2606 | | 0.0666 | | |
| 8/23/97 | 0615-0900 | 0.3878 | 0.3878 | 0.0127 | 0.0127 | 0.1400 | 0.1400 | 0.0430 | 0.0430 | |
| 8/23/97 | 0615-0900 | - | | - | | - | | - | | |
| 8/23/97 | 0903-1200 | 0.0224 | 0.0927* | 0.0407 | 0.0036* | 0.0023 | 0.0316* | 0.0026 | 0.0110* | (Mass detector stopped during analysis, sample destroyed) |
| 8/23/97 | 0903-1200 | 0.0927 | | 0.0036 | | 0.0316 | | 0.0110 | | Altered monoterpene profile |

Table B-1.1. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Azusa (continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|---------------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|---|
| 8/23/97 | 1208-1500 | 0.0127 | 0.0163 | 0.0074 | 0.0047 | 0.0021 | 0.0025 | 0.0016 | 0.0022 | |
| 8/23/97 | 1208-1500 | 0.0199 | 0.0019 | | | 0.0030 | | 0.0029 | | |
| 8/23/97 | 1505-1800 | 0.0038* | 0.0188 | 0.0188* | 0.0014 | 0.0014* | | ** | ** | |
| 8/23/97 | 1505-1800 | 0.0008 | 0.0376 | | | 0.0029 | | ** | ** | Sampling tube contents transferred from broken tube |
| AZUSA | | | | | | | | | | |
| Sep 4-7, 1997 | | | | | | | | | | |
| 9/4/97 | 0300-0600 | 0.6494 | 0.5268 | 0.0135 | 0.0115 | 0.3688 | 0.2610 | 0.0919 | 0.0744 | |
| 9/4/97 | 0300-0600 | 0.4042 | 0.0094 | | | 0.1532 | | 0.0570 | | |
| 9/4/97 | 0606-0900 | 0.2020 | 0.2141 | 0.0068 | 0.0058 | 0.0063 | 0.0344 | 0.0197 | 0.0234 | |
| 9/4/97 | 0606-0900 | 0.2262 | 0.0047 | | | 0.0625 | | 0.0271 | | |
| 9/4/97 | 0904-1200 | 0.0924 | 0.0910 | 0.0032 | 0.0028 | 0.0165 | 0.0218 | 0.0110 | 0.0113 | |
| 9/4/97 | 0904-1200 | 0.0897 | 0.0024 | | | 0.0271 | | 0.0115 | | |
| 9/4/97 | 1300-1600 | 0.0195 | 0.0208 | 0.0019 | 0.0060 | 0.0035 | 0.0058 | 0.0038 | 0.0058 | |
| 9/4/97 | 1300-1600 | 0.0221 | 0.0101 | | | 0.0081 | | 0.0079 | | |
| 9/4/97 | 1700-2000 | 0.0333 | 0.0416 | 0.0190 | 0.0117 | 0.0132 | 0.0087 | 0.0100 | 0.0104 | |
| 9/4/97 | 1700-2000 | 0.0498 | 0.0043 | | | 0.0042 | | 0.0108 | | |
| 9/4/97 | 2014-2400 | 0.1401 | 0.1286 | 0.0060 | 0.0043 | 0.0211 | 0.0351 | 0.0101 | 0.0106 | |
| 9/4/97 | 2014-2400 | 0.1171 | 0.0027 | | | 0.0491 | | 0.0112 | | |
| 9/5/97 | 0006-0300 | 0.1137 | 0.1145 | 0.0155 | 0.0096 | 0.0056 | 0.0103 | 0.0065 | 0.0084 | |
| 9/5/97 | 0006-0300 | 0.1154 | 0.0037 | | | 0.0150 | | 0.0102 | | |
| 9/5/97 | 0305-0600 | 0.2745 | 0.2838 | 0.0054 | 0.0052 | 0.1250 | 0.1523 | 0.0287 | 0.0302 | |
| 9/5/97 | 0305-0600 | 0.2931 | 0.0051 | | | 0.1796 | | 0.0317 | | |
| 9/5/97 | 0615-0900 | - | 0.3920* | - | 0.0099* | - | 0.1082* | - | 0.0383* | (High toluene, analysis in the SCAN mode) |
| 9/5/97 | 0615-0900 | 0.3920 | | 0.0099 | | 0.1082 | | 0.0385 | | |
| 9/5/97 | 0903-1200 | 0.1164 | 0.1251 | 0.0066 | 0.0065 | 0.0072 | 0.0288 | 0.0117 | 0.0160 | |
| 9/5/97 | 0903-1200 | 0.1337 | | 0.0063 | | 0.0504 | | 0.0202 | | |

Table B-1.1. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Azusa
(continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|--------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|----------------------------|
| 9/5/97 | 1300-1600 | 0.0183 | 0.0178 | 0.0027 | 0.0022 | 0.0019 | 0.0032 | 0.0033 | 0.0034 | |
| 9/5/97 | 1300-1600 | 0.0174 | 0.0016 | 0.0044 | | 0.0044 | | 0.0036 | | |
| 9/5/97 | 1700-2000 | 0.0376 | 0.0356 | 0.0022 | 0.0028 | 0.0034 | 0.0028 | 0.0065 | 0.0056 | |
| 9/5/97 | 1700-2000 | 0.0336 | | 0.0034 | | 0.0023 | | 0.0048 | | |
| 9/5/97 | 2006-2400 | 0.0989 | 0.1001 | 0.0037 | 0.0043 | 0.0396 | 0.0314 | 0.0159 | 0.0153 | |
| 9/5/97 | 2006-2400 | 0.1013 | | 0.0048 | | 0.0231 | | 0.0147 | | |
| 9/6/97 | 0008-0300 | 0.2759 | 0.2821 | 0.0084 | 0.0069 | 0.0819 | 0.1693 | 0.0176 | 0.0211 | |
| 9/6/97 | 0008-0300 | 0.2883 | | 0.0053 | | 0.2567 | | 0.0245 | | |
| 9/6/97 | 0306-0600 | 0.2541 | 0.2507 | 0.0056 | 0.0065 | 0.0476 | 0.0315 | 0.0324 | 0.0302 | |
| 9/6/97 | 0306-0600 | 0.2474 | | 0.0074 | | 0.0153 | | 0.0281 | | |
| 9/6/97 | 0615-0900 | 0.3310 | 0.3292 | 0.0110 | 0.0111 | 0.0110 | 0.0098 | 0.0382 | 0.0358 | |
| 9/6/97 | 0615-0900 | 0.3273 | | 0.0113 | | 0.0086 | | 0.0334 | | |
| 9/6/97 | 0903-1200 | 0.1008 | 0.1012 | 0.0035 | 0.0039 | 0.0291 | 0.0244 | 0.0133 | 0.0132 | |
| 9/6/97 | 0903-1200 | 0.1016 | | 0.0043 | | 0.0197 | | 0.0131 | | |
| 9/6/97 | 1300-1600 | 0.0173 | 0.0190 | 0.0054 | 0.0035 | 0.0014 | 0.0025 | 0.0019 | 0.0027 | |
| 9/6/97 | 1300-1600 | 0.0208 | | 0.0016 | | 0.0035 | | 0.0036 | | |
| 9/6/97 | 1700-1959 | 0.0269 | 0.0218 | 0.0018 | 0.0027 | 0.0092 | 0.0068 | 0.0065 | 0.0048 | |
| 9/6/97 | 1700-1959 | 0.0167 | | 0.0036 | | 0.0044 | | 0.0030 | | |
| 9/6/97 | 2010-2400 | 0.0381 | 0.0419 | 0.0031 | 0.0028 | 0.0035 | 0.0151 | 0.0055 | 0.0067 | |
| 9/6/97 | 2010-2400 | 0.0457 | | 0.0025 | | 0.0267 | | 0.0080 | | |
| 9/7/97 | 0005-0300 | 0.2871 | 0.2871* | 0.0090 | 0.0090* | 0.0358 | 0.0358* | 0.0151 | 0.0151* | |
| 9/7/97 | 0005-0300 | 0.1465 | | 0.1006 | | 0.0011 | | 0.0028 | | |
| 9/7/97 | 0306-0600 | 0.4483 | 0.4648 | 0.0095 | 0.0099 | 0.2798 | 0.3051 | 0.0407 | 0.0428 | |
| 9/7/97 | 0306-0600 | 0.4814 | | 0.0104 | | 0.3304 | | 0.0449 | | |
| 9/7/97 | 0604-0900 | 0.4308 | 0.3851 | 0.0098 | 0.0105 | 0.2207 | 0.1384 | 0.0405 | 0.0330 | |
| 9/7/97 | 0604-0900 | 0.3394 | | 0.0111 | | 0.0561 | | 0.0256 | | |

Altered monoterpene profile

Table B-1.1. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Azusa (continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|-----------------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|----------------------------|
| AZUSA | | | | | | | | | | |
| SEP 28-29, 1997 | 0301-0600 | 0.2548 | 0.2524 | 0.0239 | 0.0242 | 0.0259 | 0.0256 | 0.0571 | 0.0543 | |
| 9/28/97 | 0301-0600 | 0.2501 | 0.0245 | 0.0253 | 0.0253 | 0.0515 | | | | |
| 9/28/97 | 0609-0900 | 0.2856 | 0.2744 | 0.0103 | 0.0098 | 0.2410 | 0.2305 | 0.0768 | 0.0730 | |
| 9/28/97 | 0609-0900 | 0.2632 | 0.0094 | 0.0220 | 0.0220 | 0.0692 | | | | |
| 9/28/97 | 0904-1200 | 0.0724 | 0.0687 | 0.0060 | 0.0048 | 0.0161 | 0.0234 | 0.0121 | 0.0121 | |
| 9/28/97 | 0904-1200 | 0.0651 | 0.0336 | 0.0307 | 0.0307 | 0.0121 | | | | |
| 9/28/97 | 1300-1600 | 0.0249 | 0.0236 | 0.0017 | 0.0020 | 0.0178 | 0.0150 | 0.0087 | 0.0075 | |
| 9/28/97 | 1300-1600 | 0.0223 | 0.0023 | 0.0123 | 0.0123 | 0.0064 | | | | |
| 9/28/97 | 1700-2000 | 0.0147 | 0.0160 | 0.0050 | 0.0038 | 0.0026 | 0.0067 | 0.0054 | 0.0071 | |
| 9/28/97 | 1700-2000 | 0.0173 | 0.0026 | 0.0107 | 0.0107 | 0.0088 | | | | |
| 9/28-29/97 | 2015-0300 | 0.1072 | 0.0973 | 0.0043 | 0.0061 | 0.0898 | 0.0609 | 0.0214 | 0.0170 | |
| 9/28-29/97 | 2015-0300 | 0.0875 | 0.0080 | 0.0320 | 0.0320 | 0.0125 | | | | |
| 9/29/97 | 0310-0600 | 0.2337 | 0.2222 | 0.0118 | 0.0108 | 0.1762 | 0.1888 | 0.0546 | 0.0512 | |
| 9/29/97 | 0310-0600 | 0.2108 | 0.0097 | 0.2014 | 0.2014 | 0.0478 | | | | |
| 9/29/97 | 0616-0858 | 0.1701 | 0.1784 | 0.0477 | 0.0445 | 0.0092 | 0.0122 | 0.0131 | 0.0146 | |
| 9/29/97 | 0616-0858 | 0.1868 | 0.0413 | 0.0152 | 0.0152 | 0.0160 | | | | |
| 9/29/97 | 0900-1200 | 0.0963 | 0.0964 | 0.0044 | 0.0039 | 0.0686 | 0.0858 | 0.0179 | 0.0196 | |
| 9/29/97 | 0900-1200 | 0.0966 | 0.0034 | 0.1029 | 0.1029 | 0.0212 | | | | |
| 9/29/97 | 1300-1600 | 0.0285 | 0.0262 | 0.0018 | 0.0021 | 0.0262 | 0.0212 | 0.0100 | 0.0087 | |
| 9/29/97 | 1300-1600 | 0.0239 | 0.0023 | 0.0161 | 0.0161 | 0.0074 | | | | |
| 9/29/97 | 1700-2000 | 0.0050 | 0.0086 | 0.0233 | 0.0265 | 0.0067 | 0.0042 | 0.0028 | 0.0027 | |
| 9/29/97 | 1700-2000 | 0.0122 | 0.0297 | 0.0018 | 0.0018 | 0.0026 | | | | |

Table B-1.1. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Azusa
(continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|--------------|----------------------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|----------------------------|
| AZUSA | OCT 3-4, 1997 | | | | | | | | | |
| 10/3/97 | 0301-0600 | 0.0341 | 0.0409 | 0.0075 | 0.0059 | 0.0063 | 0.0164 | 0.0045 | 0.0075 | |
| 10/3/97 | 0301-0600 | 0.0478 | 0.0043 | 0.0265 | | | | 0.0104 | | |
| 10/3/97 | 0614-0900 | 0.0567 | 0.0050 | 0.0378 | 0.0423 | 0.0134 | | 0.0135 | | |
| 10/3/97 | 0614-0900 | 0.0584 | 0.0053 | 0.0469 | | | | 0.0135 | | |
| 10/3/97 | 0904-1200 | 0.0342 | 0.0331 | 0.0053 | 0.0168 | 0.0119 | | 0.0068 | 0.0066 | |
| 10/3/97 | 0904-1200 | 0.0320 | 0.0060 | 0.0071 | | | | 0.0065 | | |
| 10/3/97 | 1300-1600 | 0.0214 | 0.0216 | 0.0049 | 0.0047 | 0.0051 | 0.0060 | 0.0060 | 0.0063 | |
| 10/3/97 | 1300-1600 | 0.0219 | 0.0044 | 0.0068 | | | | 0.0065 | | |
| 10/3/97 | 1700-2000 | 0.0163 | 0.0162 | 0.0038 | 0.0041 | 0.0067 | 0.0100 | 0.0102 | 0.0103 | |
| 10/3/97 | 1700-2000 | 0.0160 | 0.0044 | 0.0133 | | | | 0.0104 | | |
| 10/3-4/97 | 2011-0300 | 0.1076 | 0.0934 | 0.0049 | 0.0045 | 0.0584 | 0.0450 | 0.0450 | 0.0109 | 0.0095 |
| 10/3-4/97 | 2011-0300 | 0.0791 | 0.0040 | 0.0317 | | | | 0.0081 | | |
| 10/4/97 | 0304-0600 | 0.0306 | 0.0228 | 0.1293 | 0.1306 | 0.0031 | 0.0062 | 0.0025 | 0.0059 | |
| 10/4/97 | 0304-0600 | 0.0151 | 0.1318 | | | 0.0093 | | 0.0093 | | |
| 10/5/97 | 0612-0900 | 0.0803 | 0.0800 | 0.0063 | 0.0068 | 0.0283 | 0.0213 | 0.0195 | 0.0185 | |
| 10/4/97 | 0612-0900 | 0.0798 | 0.0074 | 0.0142 | | | | 0.0174 | | |
| 10/4/97 | 0901-1200 | 0.0061 | 0.0108 | 0.0794 | 0.0529 | 0.0112 | 0.0065 | 0.0081 | 0.0060 | |
| 10/4/97 | 0901-1200 | 0.0154 | 0.0263 | | | 0.0017 | | 0.0039 | | |
| 10/4/97 | 1300-1600 | 0.0099 | 0.0138 | 0.0165 | 0.0096 | 0.0091 | 0.0095 | ** | 0.0063* | |
| 10/4/97 | 1300-1600 | 0.0178 | 0.0027 | | | 0.0099 | | 0.0063 | | |
| 10/4/97 | 1700-2000 | 0.0144 | 0.0140 | 0.0045 | 0.0042 | 0.0066 | 0.0069 | 0.0059 | 0.0059 | |
| 10/4/97 | 1700-2000 | 0.0136 | 0.0040 | | | 0.0071 | | 0.0059 | | |

*Used single value.

**Not detected.

Table B-1.2. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Pine Mtn. and Mt. Baldy.

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|------------------------------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|----------------------------|
| PINE MT AUG 4-6, 1997 | | | | | | | | | | |
| 8/4/97 | 1710-2000 | 0.0200 | 0.0208 | 0.0067 | 0.0069 | 0.0105 | 0.0108 | 0.0054 | 0.0055 | |
| 8/4/97 | 1710-2000 | 0.0217 | 0.0070 | 0.0045* | 0.0045* | 0.0007 | 0.0007 | 0.0055 | 0.0055 | |
| 08/4-08/5/97 | 2025-0259 | 0.0007 | 0.0007 | ** | ** | 0.0008 | 0.0008 | 0.0003 | 0.0003 | |
| 8/5/97 | 0314-0559 | 0.0008 | 0.0010 | 0.0017 | 0.0011 | 0.0009 | 0.0009 | 0.0002 | 0.0003 | |
| 8/5/97 | 0314-0559 | 0.0011 | 0.0006 | 0.0006 | 0.0006 | 0.0009 | 0.0009 | 0.0004 | 0.0004 | |
| 8/5/97 | 0617-0858 | 0.0148 | 0.0156 | 0.0018 | 0.0020 | 0.0038 | 0.0034 | 0.0051 | 0.0052 | |
| 8/5/97 | 0617-0858 | 0.0164 | 0.0021 | 0.0021 | 0.0021 | 0.0030 | 0.0030 | 0.0053 | 0.0053 | |
| 8/5/97 | 0911-1200 | 0.0496 | 0.0528 | 0.0080 | 0.0083 | 0.0277 | 0.0312 | 0.0066 | 0.0073 | |
| 8/5/97 | 0911-1200 | 0.0560 | 0.0086 | 0.0086 | 0.0086 | 0.0346 | 0.0346 | 0.0080 | 0.0080 | |
| 8/5/97 | 1302-1600 | 0.0381 | 0.0381* | 0.0107 | 0.0107* | 0.0087 | 0.0087* | 0.0071 | 0.0071* | (Sample tube missing) |
| 8/5/97 | 1302-1600 | - | - | - | - | - | - | - | - | |
| 8/5/97 | 1700-2000 | 0.0269 | 0.0277 | 0.0080 | 0.0084 | 0.0179 | 0.0162 | 0.0062 | 0.0063 | |
| 8/5/97 | 1700-2000 | 0.0285 | 0.0088 | 0.0088 | 0.0145 | 0.0145 | 0.0145 | 0.0065 | 0.0065 | |
| 8/5/97 | 2015-0255 | 0.0028 | 0.0029 | 0.0008 | 0.0009 | 0.0011 | 0.0009 | 0.0010 | 0.0011 | |
| 8/5/97 | 2015-0255 | 0.0029 | 0.0010 | 0.0010 | 0.0007 | 0.0007 | 0.0007 | 0.0011 | 0.0011 | |
| 8/6/97 | 0310-0600 | 0.0045 | 0.0045 | 0.0012 | 0.0014 | 0.0009 | 0.0012 | 0.0022 | 0.0022 | |
| 8/6/97 | 0310-0600 | 0.0045 | 0.0015 | 0.0015 | 0.0015 | 0.0015 | 0.0015 | 0.0021 | 0.0021 | |
| 8/6/97 | 0615-0900 | 0.0169 | 0.0164 | 0.0028 | 0.0059 | 0.0022 | 0.0017 | 0.0050 | 0.0039 | |
| 8/6/97 | 0615-0900 | 0.0159 | 0.0090 | 0.0090 | 0.0013 | 0.0013 | 0.0013 | 0.0028 | 0.0028 | |
| 8/6/97 | 0915-1200 | 0.0304 | 0.0313 | 0.0048 | 0.0051 | 0.0226 | 0.0179 | 0.0042 | 0.0043 | |
| 8/6/97 | 0915-1200 | 0.0322 | 0.0054 | 0.0054 | 0.0132 | 0.0132 | 0.0132 | 0.0044 | 0.0044 | |
| 8/6/97 | 1300-1600 | 0.0391 | 0.0402 | 0.0102 | 0.0106 | 0.0209 | 0.0153 | 0.0067 | 0.0069 | |
| 8/6/97 | 1300-1600 | 0.0413 | 0.0111 | 0.0098 | 0.0098 | 0.0098 | 0.0098 | 0.0070 | 0.0070 | |

Table B-1.2. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Pine Mtn. and Mt. Baldy (continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|---------|---------------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|----------------------------|
| PINE MT | SEP 4-7, 1997 | | | | | | | | | |
| 9/4/97 | 1702-2000 | 0.0096 | 0.0136 | 0.0106 | 0.0071 | 0.0087 | 0.0143 | 0.0015 | 0.0028 | |
| 9/4/97 | 1702-2000 | 0.0177 | 0.0036 | 0.0036 | 0.0006 | 0.0199 | 0.0014 | 0.0040 | | |
| 9/4/97 | 2008-2400 | 0.0006 | 0.0006 | 0.0008 | 0.0008 | 0.0019 | 0.0014 | 0.0007 | 0.0008 | |
| 9/4/97 | 2008-2400 | 0.0005 | 0.0009 | 0.0008 | 0.0009 | 0.0008 | 0.0010 | 0.0010 | | |
| 9/5/97 | 0010-0338 | 0.0006 | 0.0015 | 0.0082 | 0.0012 | 0.0012* | 0.0023 | 0.0023* | | |
| 9/5/97 | 0010-0338 | ** | 0.0149 | ** | ** | ** | ** | ** | | |
| 9/5/97 | 0349-0600 | 0.0009 | 0.0007 | 0.0012 | 0.0052 | 0.0027 | 0.0019 | 0.0006 | 0.0006* | |
| 9/5/97 | 0349-0600 | 0.0005 | 0.0093 | 0.0029 | 0.0010 | 0.0010 | 0.0010 | ** | | |
| 9/5/97 | 0615-0900 | 0.0038 | 0.0047 | 0.0252 | 0.0140 | 0.0016 | 0.0017 | 0.0014 | 0.0022 | |
| 9/5/97 | 0615-0900 | 0.0055 | 0.0029 | 0.0029 | 0.0017 | 0.0017 | 0.0017 | 0.0030 | | |
| 9/5/97 | 0908-1200 | 0.0305 | 0.0290 | 0.0035 | 0.0034 | 0.0206 | 0.0133 | 0.0025 | 0.0024 | |
| 9/5/97 | 0908-1200 | 0.0275 | 0.0033 | 0.0033 | 0.0060 | 0.0060 | 0.0023 | 0.0023 | | |
| 9/5/97 | 1300-1600 | 0.0288 | 0.0286 | 0.0049 | 0.0050 | 0.0040 | 0.0047 | 0.0033 | 0.0032 | |
| 9/5/97 | 1300-1600 | 0.0285 | 0.0050 | 0.0050 | 0.0054 | 0.0054 | 0.0031 | | | |
| 9/5/97 | 1700-2000 | 0.0271 | 0.0277 | 0.0042 | 0.0039 | 0.0086 | 0.0132 | 0.0038 | 0.0041 | |
| 9/5/97 | 1700-2000 | 0.0282 | 0.0035 | 0.0035 | 0.0177 | 0.0177 | 0.0043 | | | |
| 9/5/97 | 2007-2355 | 0.0007 | 0.0008 | ** | ** | 0.0014 | 0.0013 | ** | ** | |
| 9/5/97 | 2007-2355 | 0.0010 | ** | ** | 0.0013 | ** | ** | | | |
| 9/6/97 | 0000-0255 | 0.0007 | 0.0007 | ** | 0.0010 | 0.0010* | 0.0009 | 0.0009* | | |
| 9/6/97 | 0000-0255 | 0.0007 | ** | ** | ** | ** | ** | | | |
| 9/6/97 | 0301-0600 | 0.0009 | 0.0009 | 0.0013 | 0.0012 | 0.0007* | 0.0007* | ** | 0.0008* | |
| 9/6/97 | 0301-0600 | 0.0009 | 0.0012 | 0.0012 | 0.0007 | 0.0007 | 0.0008 | | | |
| 9/6/97 | 0610-0900 | 0.0033 | 0.0040 | 0.0013 | 0.0016 | 0.0024 | 0.0014 | 0.0015 | 0.0015 | |
| 9/6/97 | 0610-0900 | 0.0047 | 0.0020 | 0.0020 | 0.0004 | 0.0004 | 0.0014 | | | |
| 9/6/97 | 0905-1200 | 0.0053 | 0.0147 | 0.0048 | 0.0040 | 0.0016 | 0.0035 | ** | 0.0017* | |
| 9/6/97 | 0905-1200 | 0.0240 | 0.0033 | 0.0033 | 0.0055 | | | 0.0017 | | |

Table B-1.2. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Pine Mtn. and Mt. Baldy (continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|-----------------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|----------------------------|
| 9/6/97 | 1300-1600 | 0.0153 | 0.0154 | 0.0031 | 0.0035 | 0.0064 | 0.0046 | 0.0026 | 0.0023 | |
| 9/6/97 | 1300-1600 | 0.0154 | 0.0039 | 0.0028 | 0.0020 | 0.0050 | 0.0050 | 0.0025 | 0.0030 | |
| 9/6/97 | 1700-2000 | 0.0169 | 0.0176 | 0.0050 | 0.0046 | 0.0012 | 0.0050 | 0.0025 | 0.0030 | |
| 9/6/97 | 1700-2000 | 0.0182 | 0.0042 | 0.0088 | 0.0088 | 0.0035 | 0.0035 | 0.0035 | 0.0035 | |
| 9/6/97 | 2007-2355 | 0.0006 | 0.0010 | 0.010* | 0.0010* | 0.0009 | 0.0009 | 0.0013 | 0.0012 | |
| 9/6/97 | 2007-2355 | 0.0006 | ** | ** | ** | 0.0009 | 0.0009 | 0.0010 | 0.0010 | |
| 9/7/97 | 0000-0255 | 0.0008 | 0.0008 | 0.0010 | 0.0085 | 0.0033 | 0.0022 | 0.0018 | 0.0018* | |
| 9/7/97 | 0000-0255 | 0.0008 | 0.0160 | 0.0112 | 0.0112 | 0.0012 | 0.0012 | ** | ** | |
| 9/7/97 | 0300-0555 | 0.0016 | 0.0014 | 0.0014 | 0.0015 | 0.0011 | 0.0016 | 0.0014 | 0.0018 | |
| 9/7/97 | 0300-0555 | 0.0013 | 0.0015 | 0.0015 | 0.0021 | 0.0021 | 0.0022 | 0.0022 | 0.0022 | |
| 9/7/97 | 0600-0900 | 0.0046 | 0.0061 | 0.0024 | 0.0020 | 0.0011 | 0.0030 | 0.0008 | 0.0024 | |
| 9/7/97 | 0600-0900 | 0.0076 | 0.0017 | . | 0.0050 | 0.0050 | 0.0041 | 0.0041 | 0.0041 | |
| MT BALDY | | SEP 28-29, 1997 | | | | | | | | |
| 9/28/97 | 0904-1200 | 0.2032 | 0.2080 | 0.0091 | 0.0083 | 0.0141 | 0.0229 | 0.0563 | 0.0600 | |
| 9/28/97 | 0904-1200 | 0.2128 | 0.0076 | 0.0317 | 0.0317 | 0.0637 | 0.0637 | 0.0637 | 0.0637 | |
| 9/28/97 | 1300-1600 | 0.0998 | 0.1013 | 0.0049 | 0.0053 | 0.0183 | 0.0161 | 0.0233 | 0.0236 | |
| 9/28/97 | 1300-1600 | 0.1029 | 0.0058 | 0.0139 | 0.0139 | 0.0238 | 0.0238 | 0.0238 | 0.0238 | |
| 9/28/97 | 1700-2000 | 0.1346 | 0.1316 | 0.0111 | 0.0142 | 0.0109 | 0.0075 | 0.0472 | 0.0397 | |
| 9/28/97 | 1700-2000 | 0.1286 | 0.0172 | 0.0041 | 0.0041 | 0.0322 | 0.0322 | 0.0322 | 0.0322 | |
| 9/28-29/97 | 2005-0300 | 0.0526 | 0.0520 | 0.0056 | 0.0052 | 0.0045 | 0.0366 | 0.0353 | 0.0353 | |
| 9/28-29/97 | 2005-0300 | 0.0513 | 0.0057 | 0.0038 | 0.0038 | 0.0340 | 0.0340 | 0.0340 | 0.0340 | |
| 9/29/97 | 0307-0600 | 0.0590 | 0.0583 | 0.0055 | 0.0052 | 0.0061 | 0.0380 | 0.0376 | 0.0376 | |
| 9/29/97 | 0307-0600 | 0.0576 | 0.0050 | 0.0061 | 0.0061 | 0.0372 | 0.0372 | 0.0372 | 0.0372 | |
| 9/29/97 | 0606-0900 | 0.0748 | 0.0762 | 0.0057 | 0.0057 | 0.0079 | 0.0090 | 0.0418 | 0.0433 | |
| 9/29/97 | 0606-0900 | 0.0775 | 0.0056 | 0.0101 | 0.0101 | 0.0449 | 0.0449 | 0.0449 | 0.0449 | |
| 9/29/97 | 0903-1200 | 0.0827 | 0.0833 | 0.0124 | 0.0120 | 0.0036 | 0.0140 | 0.0146 | 0.0146 | |
| 9/29/97 | 0903-1200 | 0.0840 | 0.0117 | 0.0055 | 0.0055 | 0.0153 | 0.0153 | 0.0153 | 0.0153 | |

Table B-1.2. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Pine Mtn. and Mt. Baldy (continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|-----------------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|---------------------------------|
| 9/29/97 | 1300-1600 | 0.0431 | 0.0526 | 0.0096 | 0.0066 | 0.0071 | 0.0090 | 0.0057 | 0.0100 | |
| 9/29/97 | 1300-1600 | 0.0621 | | 0.0036 | | 0.0109 | | 0.0142 | | |
| 9/29/97 | 1700-2000 | 0.0837 | 0.0840 | 0.0079 | 0.0102 | 0.0105 | 0.0075 | 0.0332 | 0.0311 | |
| 9/29/97 | 1700-2000 | 0.0843 | | 0.0125 | | 0.0044 | | 0.0291 | | |
| MT BALDY | | OCT 3-4, 1997 | | | | | | | | |
| 10/3/97 | 0619-0900 | 0.0396 | 0.0397 | 0.0059 | | 0.0070 | | 0.0089 | 0.0177 | 0.0217 |
| 10/3/97 | 0619-0900 | 0.0398 | | 0.0059 | | 0.0107 | | 0.0257 | | |
| 10/3/97 | 0903-1200 | 0.0565 | 0.0498 | 0.0064 | 0.0080 | 0.0089 | 0.0074 | 0.0234 | 0.0172 | |
| 10/3/97 | 0903-1200 | 0.0431 | | 0.0096 | | 0.0060 | | 0.0111 | | |
| 10/3/97 | 1301-1600 | 0.0500* | | 0.0370 | 0.0068* | 0.0018 | 0.0113* | 0.0017 | 0.0102* | Altered monoterpene profile |
| 10/3/97 | 1301-1600 | 0.0500 | | 0.0068 | | 0.0113 | | 0.0102 | | |
| 10/3/97 | 1703-2003 | 0.0354 | 0.0354* | 0.0078 | 0.0078* | 0.0082 | 0.0082* | 0.0141 | 0.0141* | |
| 10/3/97 | 1703-2003 | ** | | ** | | ** | | ** | | Seems like a back sampling tube |
| 10/3-4/97 | 2010-0302 | 0.0198 | 0.0195 | 0.0028 | 0.0028 | 0.0030 | 0.0029 | 0.0155 | 0.0152 | |
| 10/3-4/97 | 2010-0302 | 0.0192 | | 0.0027 | | 0.0028 | | 0.0149 | | |
| 10/4/97 | 0310-0601 | 0.0250 | 0.0253 | 0.0040 | 0.0047 | 0.0028 | 0.0030 | 0.0218 | 0.0204 | |
| 10/4/97 | 0310-0601 | 0.0255 | | 0.0053 | | 0.0032 | | 0.0190 | | |
| 10/4/97 | 0612-0900 | 0.0458 | 0.0458* | 0.0054 | 0.0054* | 0.0080 | 0.0080* | 0.0325 | 0.0325* | |
| 10/4/97 | 0612-0900 | 0.0281 | | 0.0206 | | 0.0043 | | 0.0078 | | |
| 10/4/97 | 0903-1200 | 0.0660 | 0.0632 | 0.0087 | 0.0091 | 0.0033 | 0.0032 | 0.0109 | 0.0106 | |
| 10/4/97 | 0903-1200 | 0.0604 | | 0.0096 | | 0.0031 | | 0.0103 | | |
| 10/4/97 | 1300-1600 | 0.0451 | 0.0459 | 0.0056 | 0.0058 | 0.0102 | 0.0099 | 0.0107 | 0.0106 | |
| 10/4/97 | 1300-1600 | 0.0467 | | 0.0059 | | 0.0096 | | 0.0105 | | |
| 10/4/97 | 1700-2000 | 0.0390 | 0.0389 | 0.0074 | 0.0072 | 0.0048 | 0.0056 | 0.0126 | 0.0127 | |
| 10/4/97 | 1700-2000 | 0.0389 | | 0.0071 | | 0.0064 | | 0.0129 | | |

*Used single value.

**Not detected.

Table B-1.3. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Banning.

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|----------------|----------------------------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|---|
| BANNING | | | | | | | | | | |
| 8/4/97 | AUG 4-6, 1997 0600-0855 | 0.0283 | 0.0347 | 0.0028 | 0.0031 | 0.0061 | 0.0096 | 0.0054 | 0.0065 | |
| 8/4/97 | 0600-0855 | 0.0411 | 0.0035 | 0.0139 | 0.0009 | 0.0131 | 0.0075 | | | |
| 8/4/97 | 0902-1157 | 0.0129 | ** | 0.0009* | 0.0048 | 0.0074 | 0.0037 | 0.0041 | | |
| 8/4/97 | 0902-1157 | 0.0150 | | 0.0101 | | 0.0046 | | | | |
| 8/4/97 | - | - | - | - | - | - | - | - | - | Wind knocked over sampling equipment, no sample equipment |
| 8/4/97 | - | - | - | - | - | - | - | - | - | Wind knocked over sampling equipment, no sample equipment |
| 8/4/97 | 1504-1800 | 0.0113 | 0.0114 | 0.0014 | 0.0023 | 0.0047 | 0.0042 | 0.0032 | 0.0027 | |
| 8/4/97 | 1504-1800 | 0.0115 | 0.0032 | 0.0083 | 0.0014 | 0.0018 | 0.0037 | 0.0022 | | |
| 8/4-5/1997 | 1807-0255 | 0.0081 | 0.0013 | 0.0211 | 0.0020 | 0.0022 | 0.0019 | 0.0016 | 0.0026 | |
| 8/4-5/1997 | 1807-0255 | 0.0085 | 0.0014 | 0.0023 | 0.0020 | 0.0020 | 0.0020 | 0.0014 | 0.0027 | |
| 8/5/97 | 0303-0550 | 0.0205 | 0.0013 | 0.0602 | 0.0045 | 0.0047 | 0.0079 | 0.0059 | 0.0053 | 0.0055 |
| 8/5/97 | 0303-0550 | 0.0216 | 0.0022 | 0.0323 | 0.0013 | 0.0013 | 0.0018 | 0.0020 | 0.0020 | |
| 8/5/97 | 0604-0855 | 0.0588 | 0.0616 | 0.0013 | 0.0049 | 0.0047 | 0.0079 | 0.0069 | 0.0066 | |
| 8/5/97 | 0604-0855 | 0.0616 | 0.0013 | 0.0334 | 0.0014 | 0.0014 | 0.0021 | 0.0059 | 0.0056 | |
| 8/5/97 | 0901-1153 | 0.0312 | 0.0323 | 0.0299 | 0.0018 | 0.0018 | 0.0019 | 0.0019 | 0.0019 | |
| 8/5/97 | 0901-1153 | 0.0334 | 0.0013 | 0.0170 | 0.0188 | 0.0013 | 0.0018 | 0.0018 | 0.0018 | |
| 8/5/97 | 1159-1452 | 0.0299 | 0.0314 | 0.0170 | 0.0113 | 0.0013 | 0.0018 | 0.0018 | 0.0018 | |
| 8/5/97 | 1159-1452 | 0.0329 | 0.0170 | 0.0188 | 0.0014 | 0.0014 | 0.0021 | 0.0021 | 0.0021 | |
| 8/5/97 | 1458-1800 | 0.0207 | 0.0170 | 0.0188 | 0.0015 | 0.0015 | 0.0020 | 0.0020 | 0.0020 | |
| 8/5/97 | 1458-1800 | 0.0207 | 0.0170 | 0.0188 | 0.0015 | 0.0015 | 0.0021 | 0.0021 | 0.0021 | |
| 8/5-8/6/97 | 1816-0255 | 0.0082 | 0.0083 | 0.0022 | 0.0023 | 0.0014 | 0.0011 | 0.0011 | 0.0011 | 0.0055 |
| 8/5-8/6/97 | 1816-0255 | 0.0085 | 0.0025 | 0.0007 | 0.0007 | 0.0008 | 0.0008 | 0.0008 | 0.0008 | 0.0055 |
| 8/6/97 | 0301-0550 | 0.0027 | 0.0037 | 0.0015 | 0.0011 | 0.0018 | 0.0027 | 0.0027 | 0.0027 | 0.0056 |
| 8/6/97 | 0301-0550 | 0.0046 | 0.0322 | 0.0008 | 0.0008 | 0.0036 | 0.0036 | 0.0036 | 0.0036 | 0.0056 |
| 8/6/97 | 0605-0855 | 0.0336 | 0.0008 | 0.0008 | 0.0044 | 0.0032 | 0.0108 | 0.0108 | 0.0108 | 0.0118 |
| 8/6/97 | 0605-0855 | 0.0336 | | 0.0056 | | 0.0127 | | | | |

Table B-1.3. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Banning (continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|--------------------------------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|----------------------------|
| 8/6/97 | 0901-1150 | 0.0446 | 0.0445 | 0.0004 | 0.0019 | 0.0040 | 0.0033 | 0.0153 | 0.0124 | |
| 8/6/97 | 0901-1150 | 0.0444 | 0.0034 | 0.0006 | 0.0006 | 0.0055 | 0.0052 | 0.0094 | 0.0168 | |
| 8/6/97 | 1155-1455 | 0.0439 | 0.0466 | 0.0006 | 0.0006 | 0.0049 | 0.0047 | 0.0159 | 0.0177 | |
| 8/6/97 | 1155-1455 | 0.0494 | 0.0006 | 0.0014 | 0.0017 | 0.0033 | 0.0073 | 0.0073 | 0.0073 | |
| 8/6/97 | 1500-1800 | 0.0227 | 0.0243 | 0.0014 | 0.0019 | 0.0020 | 0.0074 | | | |
| 8/6/97 | 1500-1800 | 0.0259 | | | | | | | | |
| BANNING AUG 22-23, 1997 | | | | | | | | | | |
| 8/22/97 | 0600-0855 | 0.0383 | 0.0393 | 0.0030 | 0.0029 | 0.0034 | 0.0043 | 0.0051 | 0.0055 | |
| 8/22/97 | 0600-0855 | 0.0402 | 0.0027 | 0.0020 | 0.0022 | 0.0052 | 0.0059 | | | |
| 8/22/97 | 0900-1155 | 0.0181 | 0.0193 | 0.0025 | 0.0017 | 0.0018 | 0.0017 | 0.0019 | 0.0021 | |
| 8/22/97 | 0900-1155 | 0.0206 | 0.0095 | 0.0020 | 0.0021 | 0.0007 | 0.0007 | 0.0023 | | |
| 8/22/97 | 1200-1455 | 0.0089 | 0.0095 | 0.0021 | 0.0021 | 0.0007 | 0.0007 | 0.0010 | 0.0009 | |
| 8/22/97 | 1200-1455 | 0.0101 | 0.0102 | 0.0014 | 0.0016 | 0.0004 | 0.0004 | 0.0008 | | |
| 8/22/97 | 1500-1800 | 0.0099 | 0.0106 | 0.0017 | 0.0017 | 0.0016 | 0.0010 | 0.0015 | 0.0016 | |
| 8/22/97 | 1500-1800 | 0.0106 | 0.0052 | 0.0021 | 0.0020 | 0.0005 | 0.0006 | 0.0017 | | |
| 8/22-23/97 | 1813-0300 | 0.0049 | 0.0054 | 0.0019 | 0.0007 | 0.0007 | 0.0011 | 0.0010 | 0.0011 | |
| 8/22-23/97 | 1813-0300 | 0.0054 | | | | | | | | |
| 8/23/97 | 0303-0555 | 0.0007 | 0.0006 | 0.0012 | 0.0024 | 0.0041 | 0.0023 | ** | ** | |
| 8/23/97 | 0303-0555 | 0.0005 | 0.0036 | 0.0005 | | | ** | | | |
| 8/23/97 | 0603-0900 | 0.0135 | 0.0139 | 0.0022 | 0.0018 | 0.0015 | 0.0016 | 0.0017 | 0.0019 | |
| 8/23/97 | 0603-0900 | 0.0142 | 0.0014 | 0.0014 | 0.0018 | 0.0018 | 0.0021 | | | |
| 8/23/97 | 0903-1200 | 0.0067 | 0.0071 | 0.0013 | 0.0014 | 0.0009 | 0.0010 | 0.0010 | 0.0011 | |
| 8/23/97 | 0903-1200 | 0.0076 | 0.0014 | 0.0014 | 0.0011 | 0.0011 | 0.0012 | | | |
| 8/23/97 | 1202-1500 | 0.0024 | 0.0038 | 0.0075 | 0.0043 | 0.0008 | 0.0007 | 0.0004 | 0.0005 | |
| 8/23/97 | 1202-1500 | 0.0052 | 0.0011 | 0.0011 | 0.0006 | 0.0006 | 0.0005 | | | |
| 8/23/97 | 1502-1800 | 0.0048 | 0.0049 | 0.0013 | 0.0013 | 0.0004 | 0.0005 | 0.0007 | 0.0008 | |
| 8/23/97 | 1502-1800 | 0.0051 | 0.0013 | 0.0005 | 0.0005 | 0.0005 | 0.0008 | | | |

Table B-1.3. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Banning (continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|------------------------------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|----------------------------|
| BANNING SEP 5-6, 1997 | | | | | | | | | | |
| 9/5/97 | 0600-0855 | 0.0620 | 0.0639 | 0.0041 | 0.0049 | 0.0041 | 0.0042 | 0.0071 | 0.0074 | |
| 9/5/97 | 0600-0855 | 0.0657 | 0.0058 | 0.0016 | 0.0017 | 0.0016 | 0.0016 | 0.0077 | | |
| 9/5/97 | 0900-1155 | 0.0096 | 0.0101 | 0.0018 | 0.0009 | 0.0008 | 0.0014 | 0.0014 | 0.0014 | |
| 9/5/97 | 0900-1155 | 0.0107 | 0.0092 | 0.0009 | 0.0008 | 0.0014 | 0.0014 | 0.0014 | 0.0014 | |
| 9/5/97 | 1158-1458 | 0.0088 | 0.0096 | 0.0008 | 0.0008 | 0.0014 | 0.0014 | 0.0010 | 0.0010 | |
| 9/5/97 | 1158-1458 | 0.0096 | 0.0073 | 0.0075 | 0.0014 | 0.0014 | 0.0011 | 0.0012 | 0.0010 | |
| 9/5/97 | 1501-1800 | 0.0073 | 0.0077 | 0.0014 | 0.0014 | 0.0012 | 0.0012 | 0.0008 | 0.0009 | |
| 9/5/97 | 1501-1800 | 0.0077 | 0.0105* | 0.0015* | 0.0015* | 0.0009 | 0.0009* | 0.0020* | 0.0020* | |
| 9/5-6/97 | 1824-0257 | ** | ** | ** | ** | ** | ** | ** | ** | |
| 9/6/97 | 0308-0550 | 0.0172 | 0.0172* | 0.0021 | 0.0021* | 0.0018 | 0.0018* | 0.0034 | 0.0034* | |
| 9/6/97 | 0308-0550 | - | - | - | - | - | - | - | - | |
| 9/6/97 | 0610-0855 | 0.0178 | 0.0121* | 0.0009 | 0.0096* | 0.0018 | 0.0022* | 0.0045 | 0.0015* | |
| 9/6/97 | 0610-0855 | 0.0121 | 0.0096 | 0.0032 | 0.0022 | 0.0010* | 0.0010* | 0.0015 | 0.0015 | |
| 9/6/97 | 0900-1158 | 0.0104 | 0.0160 | 0.0054 | 0.0054 | -** | -** | -** | -** | |
| 9/6/97 | 0900-1158 | 0.0217 | 0.0069 | 0.0009 | 0.0010 | 0.0008 | 0.0008 | 0.0046* | 0.0046* | |
| 9/6/97 | 1202-1458 | 0.0064 | 0.0069 | 0.0008 | 0.0010 | 0.0009 | 0.0009 | 0.0007 | 0.0007 | |
| 9/6/97 | 1205-1458 | 0.0073 | 0.0069* | 0.0015 | 0.0011 | 0.0011* | 0.0011* | 0.0009 | 0.0009 | |
| 9/6/97 | 1502-1800 | 0.0069 | 0.0069* | 0.0163 | 0.0163 | 0.0074 | 0.0097 | 0.0009* | 0.0009* | |
| 9/6/97 | 1502-1800 | 0.0670 | - | - | - | - | - | - | - | Very low tricyclene |

Table B-1.3. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Banning
(continued).

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|------------------------------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|-----------------------------|
| BANNING OCT 3-4, 1997 | | | | | | | | | | |
| 10/3/97 | 0600-0858 | 0.0142 | 0.0142* | 0.0231 | 0.0231* | 0.0009 | 0.0009* | 0.0022 | 0.0022* | |
| 10/3/97 | 0600-0858 | 0.0043 | 0.1494 | -** | -** | -** | -** | -** | -** | Altered monoterpene profile |
| 10/3/97 | 0900-1158 | 0.0038 | 0.0027 | 0.0597 | 0.0732 | -** | -** | -** | -** | |
| 10/3/97 | 0900-1158 | 0.0016 | | 0.0867 | -** | -** | -** | -** | -** | |
| 10/3/97 | 1200-1458 | 0.0033 | 0.0030 | 0.0395 | 0.0490 | -** | 0.0015* | -** | -** | |
| 10/3/97 | 1200-1458 | 0.0028 | | 0.0586 | 0.0015 | -** | -** | -** | -** | |
| 10/3/97 | 1500-1800 | 0.0069 | 0.0069* | 0.0037 | 0.0037* | 0.0014 | 0.0014* | 0.0016 | 0.0016* | |
| 10/3/97 | 1500-1800 | 0.0019 | | 0.0681 | -** | -** | -** | -** | -** | Altered monoterpene profile |
| 10/3-4/97 | 1807-0258 | 0.0013 | 0.0091* | 0.0366 | 0.0096* | -** | 0.0007* | -** | 0.0016* | Altered monoterpene profile |
| 10/3-4/97 | 1807-0258 | 0.0091 | | 0.0096 | 0.0007 | -** | -** | -** | -** | Altered monoterpene profile |
| 10/4/97 | 0300-0553 | 0.1984 | 0.1922 | 0.0070 | 0.0091 | 0.0034 | 0.0037 | 0.0273 | 0.0244 | |
| 10/4/97 | 0300-0553 | 0.1860 | | 0.0111 | 0.0039 | -** | -** | -** | -** | Altered monoterpene profile |
| 10/4/97 | 0601-0858 | 0.3502 | 0.3356 | 0.0105 | 0.0118 | 0.0098 | 0.0065 | 0.0393 | 0.0354 | |
| 10/4/97 | 0601-0858 | 0.3210 | | 0.0131 | 0.0031 | -** | -** | -** | -** | |
| 10/4/97 | 0900-1158 | 0.0179 | 0.0177 | 0.0038 | 0.0034 | 0.0009 | 0.0011 | 0.0026 | 0.0033 | |
| 10/4/97 | 0900-1158 | 0.0174 | | 0.0031 | 0.0014 | -** | -** | -** | -** | |
| 10/4/97 | 1200-1458 | 0.0075 | 0.0074 | 0.0019 | 0.0019 | 0.0016 | 0.0015 | 0.0012 | 0.0012 | |
| 10/4/97 | 1200-1458 | 0.0073 | | 0.0020 | 0.0015 | -** | -** | -** | -** | |
| 10/4/97 | 1500-1800 | 0.0060 | | 0.0022 | 0.0033 | 0.0007 | 0.0008 | 0.0014 | 0.0012 | |
| 10/4/97 | 1500-1800 | 0.0060 | | 0.0044 | 0.0009 | -** | -** | -** | -** | |

*Used single value.

**Not detected.

Table B-1.4. Detailed Data from Measurements of α -Pinene, Camphene, Sabinene and β -Pinene in Ambient Air at Los Angeles, North Main Street.

| Date | Sampling | α -Pinene ppbv | α -Pinene av, ppbv | Camphene ppbv | Camphene av, ppbv | Sabinene ppbv | Sabinene av, ppbv | β -Pinene ppbv | β -Pinene av, ppbv | Reason for discarding data |
|-----------------------------------|-----------|--------------------------|------------------------------|------------------|----------------------|------------------|----------------------|-------------------------|-----------------------------|----------------------------|
| LOS ANGELES NORTH MAIN ST. | | | | | | | | | | |
| 8/22/97 | 0528-0855 | 0.1977 | 0.2916 | 0.0302 | 0.0217 | 0.0095 | 0.0503 | 0.0141 | 0.0324 | |
| 8/22/97 | 0558-0855 | 0.3855 | 0.0132 | 0.0911 | 0.0506 | | | | | |
| 8/22/97 | 0900-1155 | 0.0421 | 0.0361 | 0.0263 | 0.0273 | 0.0018 | 0.0042 | 0.0041 | 0.0033 | |
| 8/22/97 | 0900-1155 | 0.0302 | 0.0283 | 0.0065 | 0.0032 | 0.0082 | 0.012 | 0.0024 | 0.0025 | |
| 8/22/97 | 1202-1457 | 0.0057 | 0.0048 | 0.0032 | 0.0132 | 0.0035 | 0.0012 | 0.0022 | 0.0016 | |
| 8/22/97 | 1202-1457 | 0.0039 | 0.0044 | 0.0016 | 0.0015 | 0.0005* | 0.0005* | 0.0019 | 0.0018 | |
| 8/22/97 | 1500-1755 | 0.0042 | 0.0046 | 0.0014 | 0.0014 | 0.0005 | 0.0050 | 0.0041 | 0.0018 | |
| 8/22/97 | 1500-1755 | 0.0046 | 0.1224 | 0.0085 | 0.0084 | 0.0050 | 0.0050 | 0.0041 | 0.0310 | 0.0319 |
| 8/22-23/97 | 1806-0556 | 0.1158 | 0.1290 | 0.0084 | 0.0084 | 0.0031 | 0.0031 | 0.0035 | 0.0251 | 0.0201 |
| 8/22-23/97 | 1806-0556 | 0.1290 | 0.1764 | 0.1721 | 0.0098 | 0.0130 | 0.0051 | 0.0020 | 0.0152 | |
| 8/23/97 | 0605-0857 | 0.1677 | 0.0163 | 0.0163 | 0.0040 | 0.0044 | 0.0011 | 0.0010 | 0.0057 | 0.0048 |
| 8/23/97 | 0605-0857 | 0.0200 | 0.0197 | 0.0040 | 0.0048 | 0.0009 | 0.0009 | 0.0039 | 0.0008* | |
| 8/23/97 | 0900-1156 | 0.0194 | 0.0018 | 0.0015 | 0.0164 | 0.0003 | 0.0003* | 0.0008* | 0.0008* | |
| 8/23/97 | 0900-1156 | 0.0194 | 0.0022 | 0.0313 | 0.0004 | 0.0004 | 0.0004 | 0.0006 | 0.0004 | |
| 8/23/97 | 1202-1458 | 0.0015 | 0.0009 | 0.0010 | 0.0009 | 0.0009 | 0.0009 | 0.0003 | 0.0008* | |
| 8/23/97 | 1202-1458 | 0.0022 | 0.0009 | 0.0010 | 0.0006 | 0.0004 | 0.0004 | 0.0006 | 0.0004 | |
| 8/23/97 | 1500-1800 | 0.0009 | 0.0010 | 0.0009 | 0.0009 | 0.0009 | 0.0003 | 0.0003 | 0.0003 | |
| 8/23/97 | 1500-1800 | 0.0010 | | | | | | | | |

*Used single value.

**Not detected.

Table C-1.1. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Azusa.

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|---------------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|----------------------------|
| AZUSA | | | | | | | | |
| AUG 4-6, 1997 | | | | | | | | |
| 8/4/97 | 0600-0900 | 0.1827 | 0.1878 | 0.0427 | 0.0433 | 0.0235 | 0.0216 | |
| 8/4/97 | 0600-0900 | 0.1928 | 0.0919 | 0.0439 | 0.0197 | 0.0176 | | |
| 8/4/97 | 0907-1206 | 0.0865 | 0.0805 | 0.0372 | 0.0406 | 0.0176 | 0.0180 | |
| 8/4/97 | 0907-1206 | 0.0973 | 0.0954 | 0.0439 | 0.0184 | 0.0108 | 0.0108 | |
| 8/4/97 | 1306-1603 | 0.0820 | 0.0790 | 0.0231 | 0.0221 | 0.0108 | 0.0109 | |
| 8/4/97 | 1306-1603 | 0.0790 | 0.0790 | 0.0212 | 0.0109 | 0.0109 | 0.0109 | |
| 8/4/97 | 1700-2000 | 0.0939 | 0.0939 | 0.0338 | 0.0334 | 0.0078 | 0.0072 | |
| 8/4/97 | 1700-2000 | 0.0970 | 0.1901 | 0.0330 | 0.0361 | 0.0066 | 0.0125 | |
| 8/4/97 | 2006-0300 | 0.1920 | 0.1882 | 0.0370 | 0.0379 | 0.0125 | 0.0129 | |
| 8/4/97 | 2006-0300 | 0.1882 | 0.5276 | 0.0674 | 0.0764 | 0.0133 | 0.0320 | |
| 8/5/97 | 0307-0600 | 0.5682 | 0.0688 | 0.0855 | 0.0764 | 0.0320 | | |
| 8/5/97 | 0307-0600 | 0.6088 | 0.2678 | 0.1030 | 0.0999 | 0.0332 | | |
| 8/5/97 | 0618-0900 | 0.2873 | 0.2873 | 0.0967 | 0.0539 | 0.0577 | | |
| 8/5/97 | 0618-0900 | 0.2873 | 0.0961 | 0.1004 | 0.0512 | 0.0527 | 0.0307 | |
| 8/5/97 | 0907-1202 | 0.1047 | 0.1047 | 0.0543 | 0.0543 | 0.0310 | | |
| 8/5/97 | 0907-1202 | 0.1047 | 0.0808 | 0.0304 | 0.0312 | 0.0178 | 0.0188 | |
| 8/5/97 | 1300-1600 | 0.0813 | 0.0739 | 0.0320 | 0.0255 | 0.0199 | | |
| 8/5/97 | 1300-1600 | 0.0813 | 0.0961 | 0.0737 | 0.0234 | 0.0166 | 0.0165 | |
| 8/5/97 | 1701-2000 | 0.0735 | 0.0735 | 0.0276 | 0.0276 | 0.0165 | | |
| 8/5/97 | 1701-2000 | 0.0735 | 0.2385 | 0.0499 | 0.0526 | 0.0209 | 0.0224 | |
| 8/5- 8/6/97 | 2005-0300 | 0.2643 | 0.3595 | 0.0553 | 0.0255 | 0.0240 | | |
| 8/6/97 | 0306-0600 | 0.3595 | 0.3948 | 0.0705 | 0.0723 | 0.0247 | 0.0244 | |
| 8/6/97 | 0306-0600 | 0.3948 | 0.4873 | 0.0742 | 0.0242 | 0.0242 | | |
| 8/6/97 | 0616-0900 | 0.4873 | 0.4545 | 0.0828 | 0.0768 | 0.0350 | | |
| 8/6/97 | 0616-0900 | 0.4218 | 0.1286 | 0.0709 | 0.0480 | 0.0328 | | |
| 8/6/97 | 0906-1200 | 0.1286 | 0.1275 | 0.0475 | 0.0266 | 0.0273 | | |
| 8/6/97 | 0906-1200 | 0.1264 | 0.1264 | 0.0470 | 0.0280 | | | |

Table C-1.1. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Azusa (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|------------|-----------------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|---|
| 8/6/97 | 1300-1600 | 0.0943 | 0.0939 | 0.0302 | 0.0260 | 0.0166 | 0.0172 | |
| 8/6/97 | 1300-1600 | 0.0934 | | 0.0217 | | 0.0178 | | |
| 8/6/97 | 1700-2000 | 0.0564 | 0.0541 | 0.0156 | 0.0159 | 0.0066 | 0.0065 | |
| 8/6/97 | 1700-2000 | 0.0518 | | 0.0162 | | 0.0064 | | |
| AZUSA | Aug 22-23, 1997 | | | | | | | |
| 8/22/97 | 0318-0600 | 2.2970 | 2.2970* | 0.0437 | 0.0437* | 0.0381 | 0.0381* | |
| 8/22/97 | 0318-0600 | 5.8994 | | 0.1286 | | 0.1261 | | Sampling tube contents transferred from broken tube |
| 8/22/97 | 0610-0900 | 0.0060 | 0.6929* | ** | 0.0642* | ** | 0.0300* | All results low, seems like a back sampling tube |
| 8/22/97 | 0610-0900 | 0.6929 | | 0.0642 | | 0.0300 | | |
| 8/22/97 | 0905-1157 | 0.1484 | 0.1483 | 0.0478 | 0.0461 | 0.0213 | 0.0208 | |
| 8/22/97 | 0905-1157 | 0.1482 | | 0.0445 | | 0.0204 | | |
| 8/22/97 | 1210-1500 | 0.0956 | 0.1211* | 0.0113 | 0.0237* | 0.0135 | 0.0121* | Sampling tube contents transferred from broken tube |
| 8/22/97 | 1210-1500 | 0.1211 | | 0.0237 | | 0.0121 | | |
| 8/22/97 | 1504-1800 | 0.1711 | 0.1652 | 0.0261 | 0.0268 | 0.0105 | 0.0101 | |
| 8/22/97 | 1504-1800 | 0.1593 | | 0.0276 | | 0.0097 | | |
| 8/22-23/97 | 1815-0300 | 0.1990 | 0.1986 | 0.0301 | 0.0298 | 0.0127 | 0.0123 | |
| 8/22-23/97 | 1815-0300 | 0.1982 | | 0.0295 | | 0.0120 | | |
| 8/23/97 | 0304-0600 | 0.4643 | 0.4749* | 0.0426 | 0.0511* | 0.0259 | 0.0192* | Altered monoterpene profile |
| 8/23/97 | 0304-0600 | 0.4749 | | 0.0511 | | 0.0192 | | |
| 8/23/97 | 0615-0900 | 0.4649 | 0.4649* | 0.0408* | 0.0408* | 0.0241 | 0.0241* | |
| 8/23/97 | 0615-0900 | - | | - | | - | | (Mass detector stopped during analysis, sample destroyed) |
| 8/23/97 | 0903-1200 | 0.0918 | 0.0796* | 0.0335 | 0.0316* | 0.0173 | 0.0156* | Altered monoterpene profile |
| 8/23/97 | 0903-1200 | 0.0796 | | 0.0316 | | 0.0156 | | |
| 8/23/97 | 1208-1500 | 0.0474 | 0.0435 | 0.0184 | 0.0196 | 0.0095 | 0.0098 | |
| 8/23/97 | 1208-1500 | 0.0396 | | 0.0209 | | 0.0102 | | |

Table C-1.1. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Azusa (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|---------------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|---|
| 8/23/97 | 1505-1800 | 0.0302 | 0.0302* | 0.0114 | 0.0114* | 0.0054 | 0.0054* | |
| 8/23/97 | 1505-1800 | 0.0161 | | ** | | ** | | |
| | | | | | | | | Sampling tube contents transferred from broken tube |
| AZUSA | | | | | | | | |
| Sep 4-7, 1997 | | | | | | | | |
| 9/4/97 | 0300-0600 | 0.7931 | 0.6558 | 0.0305 | 0.0257 | 0.0212 | 0.0173 | |
| 9/4/97 | 0300-0600 | 0.4786 | | 0.0209 | | 0.0134 | | |
| 9/4/97 | 0606-0900 | 0.2756 | 0.2910 | 0.0324 | 0.0308 | 0.0350 | 0.0359 | |
| 9/4/97 | 0606-0900 | 0.3064 | | 0.0292 | | 0.0368 | | |
| 9/4/97 | 0904-1200 | 0.1199 | 0.1215 | 0.0289 | 0.0296 | 0.0288 | 0.0281 | |
| 9/4/97 | 0904-1200 | 0.1231 | | 0.0303 | | 0.0275 | | |
| 9/4/97 | 1300-1600 | 0.0937 | 0.1065 | 0.0350 | 0.0346 | 0.0206 | 0.0213 | |
| 9/4/97 | 1300-1600 | 0.1194 | | 0.0342 | | 0.0220 | | |
| 9/4/97 | 1700-2000 | 0.3697 | 0.2779 | 0.0317 | 0.0317* | 0.0136 | 0.0106 | |
| 9/4/97 | 1700-2000 | 0.1861 | | ** | | 0.0076 | | |
| 9/4/97 | 2014-2400 | 0.1700 | 0.1533 | 0.0134 | 0.0152 | 0.0106 | 0.0094 | |
| 9/4/97 | 2014-2400 | 0.1365 | | 0.0169 | | 0.0081 | | |
| 9/5/97 | 0006-0300 | 0.1278 | 0.1135 | 0.0227 | 0.0207 | 0.0115 | 0.0105 | |
| 9/5/97 | 0006-0300 | 0.0991 | | 0.0186 | | 0.0094 | | |
| 9/5/97 | 0305-0600 | 0.2764 | 0.2849 | 0.0277 | 0.0292 | 0.0187 | 0.0190 | |
| 9/5/97 | 0305-0600 | 0.2934 | | 0.0306 | | 0.0192 | | |
| 9/5/97 | 0615-0900 | - | 0.4983* | - | 0.0341* | - | 0.0228* | (High toluene, analysis in the SCAN mode) |
| 9/5/97 | 0615-0900 | 0.4983 | | 0.0341 | | 0.0228 | | |
| 9/5/97 | 0903-1200 | 0.1416 | 0.1626 | 0.0323 | 0.0313 | 0.0207 | 0.0232 | |
| 9/5/97 | 0903-1200 | 0.1837 | | 0.0302 | | 0.0258 | | |
| 9/5/97 | 1300-1600 | 0.1265 | 0.1287 | 0.0180 | 0.0162 | 0.0065 | 0.0073 | |
| 9/5/97 | 1300-1600 | 0.1310 | | 0.0144 | | 0.0081 | | |
| 9/5/97 | 1700-2000 | 0.0591 | 0.0587 | 0.0096 | 0.0097 | 0.0040 | 0.0037 | |
| 9/5/97 | 1700-2000 | 0.0584 | | 0.0097 | | 0.0034 | | |

Table C-1.1. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Azusa (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|------------------------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|----------------------------|
| 9/5/97 | 2006-2400 | 0.1797 | 0.1799 | 0.0162 | 0.0146 | 0.0071 | 0.0074 | |
| 9/5/97 | 2006-2400 | 0.1800 | | 0.0130 | | 0.0078 | | |
| 9/6/97 | 0008-0300 | 0.2756 | 0.2827 | 0.0177 | 0.0190 | 0.0134 | 0.0128 | |
| 9/6/97 | 0008-0300 | 0.2899 | | 0.0203 | | 0.0122 | | |
| 9/6/97 | 0306-0600 | 0.2602 | 0.2534 | 0.0186 | 0.0186 | 0.0166 | 0.0164 | |
| 9/6/97 | 0306-0600 | 0.2465 | | 0.0186 | | 0.0162 | | |
| 9/6/97 | 0615-0900 | 0.2859 | 0.2745 | 0.0347 | 0.0350 | 0.0263 | 0.0257 | |
| 9/6/97 | 0615-0900 | 0.2630 | | 0.0354 | | 0.0250 | | |
| 9/6/97 | 0903-1200 | 0.0616 | 0.0645 | 0.0318 | 0.0333 | 0.0235 | 0.0232 | |
| 9/6/97 | 0903-1200 | 0.0673 | | 0.0348 | | 0.0229 | | |
| 9/6/97 | 1300-1600 | 0.1057 | 0.1035 | 0.0126 | 0.0130 | 0.0082 | 0.0074 | |
| 9/6/97 | 1300-1600 | 0.1013 | | 0.0134 | | 0.0066 | | |
| 9/6/97 | 1700-1959 | 0.1018 | 0.0970 | 0.0072 | 0.0077 | 0.0022 | 0.0023 | |
| 9/6/97 | 1700-1959 | 0.0922 | | 0.0082 | | 0.0024 | | |
| 9/6/97 | 2010-2400 | 0.0826 | 0.0968 | 0.0097 | 0.0095 | 0.0066 | 0.0069 | |
| 9/6/97 | 2010-2400 | 0.1110 | | 0.0093 | | 0.0072 | | |
| 9/7/97 | 0005-0300 | 0.2741 | 0.2741* | 0.0201 | 0.0201* | 0.0148 | 0.0148* | |
| 9/7/97 | 0005-0300 | 0.3651 | | 0.0200 | | 0.0266 | | |
| 9/7/97 | 0306-0600 | 0.6203 | 0.6366 | 0.0355 | 0.0362 | 0.0290 | 0.0311 | |
| 9/7/97 | 0306-0600 | 0.6529 | | 0.0369 | | 0.0332 | | |
| 9/7/97 | 0604-0900 | 0.7725 | 0.6723 | 0.0514 | 0.0454 | 0.0443 | 0.0379 | |
| 9/7/97 | 0604-0900 | 0.5721 | | 0.0394 | | 0.0315 | | |
| AZUSA | | | | | | | | |
| SEP 28-29, 1997 | | | | | | | | |
| 9/28/97 | 0301-0600 | 0.4213 | 0.4025 | 0.0796 | 0.0767 | 0.0383 | 0.0368 | |
| 9/28/97 | 0301-0600 | 0.3836 | | 0.0738 | | 0.0353 | | |
| 9/28/97 | 0609-0900 | 3.9707 | 3.7978 | 0.0650 | 0.0626 | 0.0376 | 0.0346 | |
| 9/28/97 | 0609-0900 | 3.6249 | | 0.0602 | | 0.0316 | | |
| 9/28/97 | 0904-1200 | 0.1039 | 0.0813 | 0.0436 | 0.0417 | 0.0369 | 0.0325 | |
| 9/28/97 | 0904-1200 | 0.0587 | | 0.0398 | | 0.0281 | | |

Table C-1.1. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Azusa (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|----------------------------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|----------------------------|
| 9/28/97 | 1300-1600 | 0.0620 | 0.0610 | 0.0443 | 0.0379 | 0.0279 | 0.0257 | |
| 9/28/97 | 1300-1600 | 0.0599 | | 0.0315 | | 0.0236 | | |
| 9/28/97 | 1700-2000 | 0.0663 | 0.0562 | 0.0281 | 0.0266 | 0.0190 | 0.0160 | |
| 9/28/97 | 1700-2000 | 0.0461 | | 0.0252 | | 0.0129 | | |
| 9/28-29/97 | 2015-0300 | 0.1319 | 0.1230 | 0.0306 | 0.0277 | 0.0156 | 0.0143 | |
| 9/28-29/97 | 2015-0300 | 0.1141 | | 0.0249 | | 0.0131 | | |
| 9/29/97 | 0310-0600 | 0.6488 | 0.5995 | 0.0698 | 0.0632 | 0.0463 | 0.0424 | |
| 9/29/97 | 0310-0600 | 0.5501 | | 0.0567 | | 0.0385 | | |
| 9/29/97 | 0616-0858 | 0.6943 | 0.6689 | 0.0492 | 0.0443 | 0.0331 | 0.0296 | |
| 9/29/97 | 0616-0858 | 0.6436 | | 0.0395 | | 0.0261 | | |
| 9/29/97 | 0900-1200 | 0.0979 | 0.1155 | 0.0489 | 0.0581 | 0.0238 | 0.0295 | |
| 9/29/97 | 0900-1200 | 0.1330 | | 0.0674 | | 0.0352 | | |
| 9/29/97 | 1300-1600 | 0.1915 | 0.1950 | 0.0406 | 0.0419 | 0.0222 | 0.0261 | |
| 9/29/97 | 1300-1600 | 0.1984 | | 0.0431 | | 0.0301 | | |
| 9/29/97 | 1700-2000 | 0.2560 | 0.1952 | 0.0310 | 0.0291 | 0.0237 | 0.0207 | |
| 9/29/97 | 1700-2000 | 0.1344 | | 0.0273 | | 0.0177 | | |
| AZUSA OCT 3-4, 1997 | | | | | | | | |
| 10/3/97 | 0301-0600 | 0.0831 | 0.1046 | 0.0149 | 0.0215 | 0.0126 | 0.0175 | |
| 10/3/97 | 0301-0600 | 0.1261 | | 0.0281 | | 0.0224 | | |
| 10/3/97 | 0614-0900 | 0.1566 | 0.1525 | 0.0587 | 0.0563 | 0.1143 | 0.1136 | |
| 10/3/97 | 0614-0900 | 0.1484 | | 0.0539 | | 0.1129 | | |
| 10/3/97 | 0904-1200 | 0.0769 | 0.0976 | 0.0314 | 0.0380 | 0.0639 | 0.0675 | |
| 10/3/97 | 0904-1200 | 0.1183 | | 0.0446 | | 0.0712 | | |
| 10/3/97 | 1300-1600 | 0.0899 | 0.0909 | 0.0407 | 0.0418 | 0.0246 | 0.0240 | |
| 10/3/97 | 1300-1600 | 0.0919 | | 0.0429 | | 0.0234 | | |
| 10/3/97 | 1700-2000 | 0.1081 | 0.1050 | 0.0260 | 0.0249 | 0.0245 | 0.0240 | |
| 10/3/97 | 1700-2000 | 0.1019 | | 0.0238 | | 0.0236 | | |
| 10/3-4/97 | 2011-0300 | 0.0807 | 0.0670 | 0.0251 | 0.0206 | 0.0112 | 0.0090 | |
| 10/3-4/97 | 2011-0300 | 0.0533 | | 0.0161 | | 0.0067 | | |

Table C-1.1. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Azusa (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|---------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|----------------------------|
| 10/4/97 | 0304-0600 | 0.1573 | 0.1224 | 0.0249 | 0.0249 | 0.0372 | 0.0413 | |
| 10/4/97 | 0304-0600 | 0.0875 | | 0.0248 | | 0.0454 | | |
| 10/5/97 | 0612-0900 | 0.1392 | 0.1371 | 0.0603 | 0.0606 | 0.0523 | 0.0519 | |
| 10/4/97 | 0612-0900 | 0.1349 | | 0.0609 | | 0.0515 | | |
| 10/4/97 | 0901-1200 | 0.1310 | 0.1251 | 0.0402 | 0.0393 | 0.0618 | 0.0493 | |
| 10/4/97 | 0901-1200 | 0.1192 | | 0.0384 | | 0.0367 | | |
| 10/4/97 | 1300-1600 | 0.0475 | 0.0492 | 0.0120 | 0.0214 | 0.0106 | 0.0138 | |
| 10/4/97 | 1300-1600 | 0.0509 | | 0.0308 | | 0.0171 | | |
| 10/4/97 | 1700-2000 | 0.0866 | 0.0870 | 0.0155 | 0.0159 | 0.0153 | 0.0161 | |
| 10/4/97 | 1700-2000 | 0.0874 | | 0.0162 | | 0.0168 | | |

*Used single value.

**Not detected.

Table C-1.2. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Pine Mtn. and Mt. Baldy.

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|------------------------------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|----------------------------|
| PINE MT AUG 4-6, 1997 | | | | | | | | |
| 8/4/97 | 1710-2000 | 0.0115 | 0.0107 | 0.0232 | 0.0240 | 0.0246 | 0.0246 | |
| 8/4/97 | 1710-2000 | 0.0098 | | 0.0248 | | 0.0245 | | |
| 08/4-08/5/97 | 2025-0259 | 0.0066 | 0.0050 | 0.0110 | 0.0147 | 0.0205 | 0.0214 | |
| 08/4-08/5/97 | 2025-0259 | 0.0035 | | 0.0184 | | 0.0224 | | |
| 8/5/97 | 0314-0559 | 0.0020 | 0.0021 | 0.0058 | 0.0060 | 0.0078 | 0.0079 | |
| 8/5/97 | 0314-0559 | 0.0021 | | 0.0063 | | 0.0081 | | |
| 8/5/97 | 0617-0858 | 0.0106 | 0.0111 | 0.0044 | 0.0048 | 0.0042 | 0.0043 | |
| 8/5/97 | 0617-0858 | 0.0116 | | 0.0051 | | 0.0043 | | |
| 8/5/97 | 0911-1200 | 0.0119 | 0.0124 | 0.0299 | 0.0310 | 0.0298 | 0.0299 | |
| 8/5/97 | 0911-1200 | 0.0129 | | 0.0322 | | 0.0300 | | |
| 8/5/97 | 1302-1600 | 0.0118 | 0.0118* | 0.0468 | 0.0468* | 0.0547 | 0.0547* | (Sample tube missing) |
| 8/5/97 | 1302-1600 | - | | - | | - | | |
| 8/5/97 | 1700-2000 | 0.0073 | 0.0082 | 0.0313 | 0.0329 | 0.0459 | 0.0483 | |
| 8/5/97 | 1700-2000 | 0.0090 | | 0.0344 | | 0.0506 | | |
| 8/5/97 | 2015-0255 | 0.0028 | 0.0030 | 0.0154 | 0.0157 | 0.0277 | 0.0286 | |
| 8/5/97 | 2015-0255 | 0.0031 | | 0.0160 | | 0.0295 | | |
| 8/6/97 | 0310-0600 | 0.0035 | 0.0039 | 0.0029 | 0.0036 | 0.0055 | 0.0059 | |
| 8/6/97 | 0310-0600 | 0.0043 | | 0.0042 | | 0.0063 | | |
| 8/6/97 | 0615-0900 | 0.0073 | 0.0083 | 0.0074 | 0.0070 | 0.0096 | 0.0095 | |
| 8/6/97 | 0615-0900 | 0.0093 | | 0.0065 | | 0.0095 | | |
| 8/6/97 | 0915-1200 | 0.0087 | 0.0090 | 0.0214 | 0.0227 | 0.0265 | 0.0278 | |
| 8/6/97 | 0915-1200 | 0.0094 | | 0.0240 | | 0.0290 | | |
| 8/6/97 | 1300-1600 | 0.0103 | 0.0106 | 0.0536 | 0.0563 | 0.0578 | 0.0577 | |
| 8/6/97 | 1300-1600 | 0.0109 | | 0.0589 | | 0.0575 | | |

Table C-1.2. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Pine Mtn. and Mt. Baldy (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|----------------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|----------------------------|
| PINE MT | | | | | | | | |
| SEP 4-7, 1997 | | | | | | | | |
| 9/4/97 | 1702-2000 | 0.0116 | 0.0121 | 0.0079 | 0.0092 | 0.0106 | 0.0106 | |
| 9/4/97 | 1702-2000 | 0.0126 | 0.0105 | 0.0076 | 0.0079 | 0.0106 | 0.0106 | |
| 9/4/97 | 2008-2400 | 0.0033 | 0.0033 | 0.0082 | 0.0082 | 0.0092 | 0.0079 | |
| 9/4/97 | 2008-2400 | 0.0034 | 0.0037 | -** | -** | 0.0066 | 0.0079 | |
| 9/5/97 | 0010-0338 | 0.0042 | 0.0033 | -** | -** | 0.0029 | 0.0029* | |
| 9/5/97 | 0010-0338 | 0.0033 | -** | -** | -** | -** | -** | |
| 9/5/97 | 0349-0600 | 0.0064 | 0.0062 | 0.0059 | 0.0059* | 0.0049 | 0.0044 | |
| 9/5/97 | 0349-0600 | 0.0059 | -** | -** | -** | 0.0040 | 0.0040 | |
| 9/5/97 | 0615-0900 | 0.0059 | 0.0049 | -** | -** | 0.0057 | 0.0052 | |
| 9/5/97 | 0615-0900 | 0.0038 | -** | -** | -** | 0.0048 | 0.0048 | |
| 9/5/97 | 0908-1200 | 0.0088 | 0.0076 | 0.0119 | 0.0088 | 0.0141 | 0.0127 | |
| 9/5/97 | 0908-1200 | 0.0063 | 0.0056 | 0.0121 | 0.0121 | 0.0112 | 0.0112 | |
| 9/5/97 | 1300-1600 | 0.0080 | 0.0079 | 0.0121 | 0.0121 | 0.0168 | 0.0153 | |
| 9/5/97 | 1300-1600 | 0.0079 | 0.0121 | 0.0121 | 0.0121 | 0.0138 | 0.0127 | |
| 9/5/97 | 1700-2000 | 0.0092 | 0.0090 | 0.0086 | 0.0089 | 0.0105 | 0.0103 | |
| 9/5/97 | 1700-2000 | 0.0088 | 0.0093 | 0.0093 | 0.0101 | 0.0101 | 0.0101 | |
| 9/5/97 | 2007-2355 | -** | 0.0062 | 0.0058 | 0.0072 | 0.0068 | 0.0068 | |
| 9/5/97 | 2007-2355 | -** | 0.0055 | 0.0055 | 0.0064 | 0.0064 | 0.0064 | |
| 9/6/97 | 0000-0255 | 0.0035 | 0.0032 | 0.0046 | 0.0046 | 0.0058 | 0.0053 | |
| 9/6/97 | 0000-0255 | 0.0029 | 0.0046 | 0.0046 | 0.0047 | 0.0047 | 0.0047 | |
| 9/6/97 | 0301-0600 | 0.0027 | 0.0023 | -** | -** | 0.0037 | 0.0037 | |
| 9/6/97 | 0301-0600 | 0.0019 | -** | -** | -** | 0.0037 | 0.0037 | |
| 9/6/97 | 0610-0900 | 0.0026 | 0.0028 | 0.0058 | 0.0058* | 0.0053 | 0.0056 | |
| 9/6/97 | 0610-0900 | 0.0030 | -** | -** | -** | 0.0058 | 0.0058 | |
| 9/6/97 | 0905-1200 | -** | 0.0061* | 0.0068 | 0.0090 | -** | 0.0105* | |
| 9/6/97 | 0905-1200 | 0.0061 | 0.0112 | 0.0112 | 0.0105 | 0.0105 | 0.0105 | |

Table C-1.2. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Pine Mtn. and Mt. Baldy (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|---------------------------------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|----------------------------|
| 9/6/97 | 1300-1600 | 0.0051 | 0.0048 | 0.0110 | 0.0113 | 0.0119 | 0.0106 | |
| 9/6/97 | 1300-1600 | 0.0046 | | 0.0116 | | 0.0094 | | |
| 9/6/97 | 1700-2000 | 0.0060 | 0.0057 | 0.0082 | 0.0090 | 0.0096 | 0.0096 | |
| 9/6/97 | 1700-2000 | 0.0053 | | 0.0099 | | 0.0097 | | |
| 9/6/97 | 2007-2355 | 0.0026 | 0.0025 | 0.0050 | 0.0038 | 0.0051 | 0.0046 | |
| 9/6/97 | 2007-2355 | 0.0025 | | 0.0026 | | 0.0042 | | |
| 9/7/97 | 0000-0255 | 0.0022 | 0.0026 | ** | ** | 0.0029 | 0.0036 | |
| 9/7/97 | 0000-0255 | 0.0029 | | ** | | 0.0043 | | |
| 9/7/97 | 0300-0555 | 0.0023 | 0.0026 | 0.0088 | 0.0088* | 0.0022 | 0.0025 | |
| 9/7/97 | 0300-0555 | 0.0029 | | ** | | 0.0028 | | |
| 9/7/97 | 0600-0900 | 0.0041 | 0.0035 | ** | ** | 0.0016 | 0.0015 | |
| 9/7/97 | 0600-0900 | 0.0030 | | ** | | 0.0013 | | |
| MT BALDY SEP 28-29, 1997 | | | | | | | | |
| 9/28/97 | 0904-1200 | 0.0570 | 0.0564 | 0.0142 | 0.0142 | 0.0120 | 0.0133 | |
| 9/28/97 | 0904-1200 | 0.0558 | | 0.0142 | | 0.0146 | | |
| 9/28/97 | 1300-1600 | 0.0198 | 0.0208 | 0.0204 | 0.0222 | 0.0155 | 0.0162 | |
| 9/28/97 | 1300-1600 | 0.0218 | | 0.0239 | | 0.0170 | | |
| 9/28/97 | 1700-2000 | 0.0258 | 0.0305 | 0.0232 | 0.0224 | 0.0101 | 0.0118 | |
| 9/28/97 | 1700-2000 | 0.0351 | | 0.0215 | | 0.0135 | | |
| 9/28-29/97 | 2005-0300 | 0.0115 | 0.0110 | 0.0196 | 0.0188 | 0.0109 | 0.0104 | |
| 9/28-29/97 | 2005-0300 | 0.0106 | | 0.0180 | | 0.0099 | | |
| 9/29/97 | 0307-0600 | 0.0129 | 0.0125 | 0.0139 | 0.0140 | 0.0087 | 0.0083 | |
| 9/29/97 | 0307-0600 | 0.0121 | | 0.0140 | | 0.0079 | | |
| 9/29/97 | 0606-0900 | 0.0165 | 0.0172 | 0.0156 | 0.0160 | 0.0070 | 0.0072 | |
| 9/29/97 | 0606-0900 | 0.0178 | | 0.0164 | | 0.0074 | | |
| 9/29/97 | 0903-1200 | 0.0285 | 0.0297 | 0.0201 | 0.0204 | 0.0108 | 0.0110 | |
| 9/29/97 | 0903-1200 | 0.0308 | | 0.0208 | | 0.0111 | | |

Table C-1.2. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Pine Mtn. and Mt. Baldy (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|-----------------|----------------------------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|-----------------------------|
| 9/29/97 | 1300-1600 | 0.0196 | 0.0162 | 0.0123 | 0.0121 | 0.0080 | 0.0076 | |
| 9/29/97 | 1300-1600 | 0.0128 | | 0.0120 | | 0.0072 | | |
| 9/29/97 | 1700-2000 | 0.0150 | 0.0218 | 0.0183 | 0.0192 | 0.0082 | 0.0089 | |
| 9/29/97 | 1700-2000 | 0.0286 | | 0.0200 | | 0.0096 | | |
| MT BALDY | | | | | | | | |
| 10/3/97 | OCT 3-4, 1997 0619-0900 | 0.0190 | 0.0234 | 0.0134 | 0.0181 | 0.0168 | 0.0258 | |
| 10/3/97 | 0619-0900 | 0.0279 | | 0.0228 | | 0.0349 | | |
| 10/3/97 | 0903-1200 | 0.0296 | 0.0337 | 0.0362 | 0.0308 | 0.0478 | 0.0414 | |
| 10/3/97 | 0903-1200 | 0.0378 | | 0.0254 | | 0.0349 | | |
| 10/3/97 | 1301-1600 | 0.0183 | 0.0098* | 0.0148 | 0.0234* | 0.0031 | 0.0186* | Altered monoterpene profile |
| 10/3/97 | 1301-1600 | 0.0098 | | 0.0234 | | 0.0186 | | |
| 10/3/97 | 1703-2003 | 0.0191 | 0.0191* | 0.0432 | 0.0432* | 0.0247 | 0.0247* | |
| 10/3/97 | 1703-2003 | -** | | -** | | -** | | |
| 10/3-4/97 | 2010-0302 | 0.0046 | 0.0046 | 0.0112 | 0.0118 | 0.0112 | 0.0113 | |
| 10/3-4/97 | 2010-0302 | 0.0046 | | 0.0124 | | 0.0114 | | |
| 10/4/97 | 0310-0601 | 0.0065 | 0.0074 | 0.0117 | 0.0130 | 0.0058 | 0.0066 | |
| 10/4/97 | 0310-0601 | 0.0083 | | 0.0143 | | 0.0074 | | |
| 10/4/97 | 0612-0900 | 0.0188 | 0.0188* | 0.0164 | 0.0164* | 0.0066 | 0.0066* | |
| 10/4/97 | 0612-0900 | 0.0218 | | 0.0103 | | 0.0054 | | |
| 10/4/97 | 0903-1200 | 0.0196 | 0.0201 | 0.0154 | 0.0151 | 0.0089 | 0.0081 | |
| 10/4/97 | 0903-1200 | 0.0206 | | 0.0147 | | 0.0073 | | |
| 10/4/97 | 1300-1600 | 0.0103 | 0.0114 | 0.0262 | 0.0272 | 0.0189 | 0.0208 | |
| 10/4/97 | 1300-1600 | 0.0125 | | 0.0283 | | 0.0227 | | |
| 10/4/97 | 1700-2000 | 0.0194 | 0.0178 | 0.0381 | 0.0354 | 0.0245 | 0.0243 | |
| 10/4/97 | 1700-2000 | 0.0162 | | 0.0327 | | 0.0242 | | |

*Used single value.
**Not detected.

Table C-13. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Banning.

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|------------------------------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|---|
| BANNING AUG 4-6, 1997 | | | | | | | | |
| 8/4/97 | 0600-0855 | 0.0105 | 0.0111 | 0.0154 | 0.0198 | 0.0129 | 0.0153 | |
| 8/4/97 | 0600-0855 | 0.0118 | 0.0241 | 0.0177 | | | | |
| 8/4/97 | 0902-1157 | 0.0028 | 0.0036 | 0.0048 | 0.0051 | 0.0040 | 0.0040 | |
| 8/4/97 | 0902-1157 | 0.0043 | - | 0.0054 | - | 0.0040 | - | Wind knocked over sampling equipment, no sample |
| 8/4/97 | - | - | - | - | - | - | - | Wind knocked over sampling equipment, no sample |
| 8/4/97 | 1504-1800 | 0.0055 | 0.0070 | 0.0126 | 0.0147 | 0.0175 | 0.0179 | |
| 8/4/97 | 1504-1800 | 0.0085 | 0.0169 | 0.0183 | | | | |
| 8/4-5/1997 | 1807-0255 | 0.0062 | 0.0064 | 0.0215 | 0.0222 | 0.0083 | 0.0086 | |
| 8/4-5/1997 | 1807-0255 | 0.0066 | - | 0.0228 | - | 0.0090 | - | |
| 8/5/97 | 0303-0550 | 0.0266 | 0.0273 | 0.0178 | 0.0179 | 0.0095 | 0.0098 | |
| 8/5/97 | 0303-0550 | 0.0280 | - | 0.0180 | - | 0.0101 | - | |
| 8/5/97 | 0604-0855 | 0.0230 | 0.0227 | 0.0196 | 0.0200 | 0.0125 | 0.0134 | |
| 8/5/97 | 0604-0855 | 0.0224 | - | 0.0203 | - | 0.0143 | - | |
| 8/5/97 | 0901-1153 | 0.0050 | 0.0048 | 0.0048 | 0.0047 | 0.0070 | 0.0074 | |
| 8/5/97 | 0901-1153 | 0.0047 | - | 0.0047 | - | 0.0078 | - | |
| 8/5/97 | 1159-1452 | 0.0050 | 0.0055 | 0.0026 | 0.0026 | 0.0031 | 0.0030 | |
| 8/5/97 | 1159-1452 | 0.0060 | - | 0.0026 | - | 0.0028 | - | |
| 8/5/97 | 1458-1800 | 0.0061 | 0.0065 | 0.0045 | 0.0051 | 0.0036 | 0.0036 | |
| 8/5/97 | 1458-1800 | 0.0069 | - | 0.0057 | - | 0.0035 | - | |
| 8/5- 8/6/97 | 1816-0255 | 0.0065 | 0.0069 | 0.0337 | 0.0351 | 0.0243 | 0.0248 | |
| 8/5- 8/6/97 | 1816-0255 | 0.0074 | - | 0.0364 | - | 0.0252 | - | |
| 8/6/97 | 0301-0550 | 0.0020 | 0.0033 | 0.0066 | 0.0096 | 0.0061 | 0.0097 | |
| 8/6/97 | 0301-0550 | 0.0047 | - | 0.0126 | - | 0.0134 | - | |
| 8/6/97 | 0605-0855 | 0.0040 | 0.0045 | 0.0031 | 0.0029 | 0.0022 | 0.0022 | |
| 8/6/97 | 0605-0855 | 0.0049 | - | 0.0027 | - | 0.0022 | - | |

Table C-1.3. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Banning (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|--------------------------------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|----------------------------|
| 8/6/97 | 0901-1150 | 0.0041 | 0.0051 | -** | 0.0029* | 0.0018 | 0.0017 | |
| 8/6/97 | 0901-1150 | 0.0061 | | 0.0029 | | 0.0015 | | |
| 8/6/97 | 1155-1455 | 0.0043 | 0.0048 | 0.0024 | 0.0027 | 0.0018 | 0.0018 | |
| 8/6/97 | 1155-1455 | 0.0052 | | 0.0029 | | 0.0018 | | |
| 8/6/97 | 1500-1800 | 0.0038 | 0.0041 | 0.0120 | 0.0117 | 0.0125 | 0.0121 | |
| 8/6/97 | 1500-1800 | 0.0045 | | 0.0114 | | 0.0116 | | |
| BANNING AUG 22-23, 1997 | | | | | | | | |
| 8/22/97 | 0600-0855 | 0.0251 | 0.0271 | 0.0121 | 0.0119 | 0.0073 | 0.0067 | |
| 8/22/97 | 0600-0855 | 0.0291 | | 0.0117 | | 0.0061 | | |
| 8/22/97 | 0900-1155 | 0.0068 | 0.0084 | 0.0113 | 0.0116 | 0.0066 | 0.0067 | |
| 8/22/97 | 0900-1155 | 0.0099 | | 0.0119 | | 0.0067 | | |
| 8/22/97 | 1200-1455 | 0.0030 | 0.0035 | 0.0208 | 0.0242 | 0.0255 | 0.0307 | |
| 8/22/97 | 1200-1455 | 0.0039 | | 0.0276 | | 0.0359 | | |
| 8/22/97 | 1500-1800 | 0.0037 | 0.0037 | 0.0120 | 0.0139 | 0.0128 | 0.0132 | |
| 8/22/97 | 1500-1800 | 0.0037 | | 0.0158 | | 0.0137 | | |
| 8/22-23/97 | 1813-0300 | 0.0077 | 0.0080 | 0.0295 | 0.0304 | 0.0180 | 0.0187 | |
| 8/22-23/97 | 1813-0300 | 0.0083 | | 0.0313 | | 0.0195 | | |
| 8/23/97 | 0303-0555 | 0.0020 | 0.0026 | 0.0184 | 0.0169 | 0.0169 | 0.0154 | |
| 8/23/97 | 0303-0555 | 0.0033 | | 0.0155 | | 0.0138 | | |
| 8/23/97 | 0603-0900 | 0.0029 | 0.0033 | 0.0120 | 0.0122 | 0.0108 | 0.0111 | |
| 8/23/97 | 0603-0900 | 0.0036 | | 0.0124 | | 0.0115 | | |
| 8/23/97 | 0903-1200 | 0.0019 | 0.0019 | 0.0060 | 0.0065 | 0.0064 | 0.0058 | |
| 8/23/97 | 0903-1200 | 0.0020 | | 0.0070 | | 0.0051 | | |
| 8/23/97 | 1202-1500 | 0.0021 | 0.0020 | 0.0059 | 0.0071 | 0.0061 | 0.0053 | |
| 8/23/97 | 1202-1500 | 0.0019 | | 0.0083 | | 0.0045 | | |
| 8/23/97 | 1502-1800 | 0.0019 | 0.0020 | 0.0084 | 0.0088 | 0.0103 | 0.0120 | |
| 8/23/97 | 1502-1800 | 0.0022 | | 0.0092 | | 0.0138 | | |

Table C-1.3. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Banning (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|------------------------------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|---|
| BANNING SEP 5-6, 1997 | | | | | | | | |
| 9/5/97 | 0600-0855 | 0.0101 | 0.0160 | 0.0367 | 0.0381 | 0.0125 | 0.0131 | |
| 9/5/97 | 0600-0855 | 0.0218 | | 0.0395 | | 0.0137 | | |
| 9/5/97 | 0900-1155 | 0.0026 | 0.0027 | 0.0133 | 0.0133 | 0.0048 | 0.0053 | |
| 9/5/97 | 0900-1155 | 0.0028 | | 0.0133 | | 0.0057 | | |
| 9/5/97 | 1158-1458 | 0.0028 | 0.0029 | 0.0070 | 0.0074 | 0.0053 | 0.0056 | |
| 9/5/97 | 1158-1458 | 0.0030 | | 0.0077 | | 0.0058 | | |
| 9/5/97 | 1501-1800 | 0.0032 | 0.0033 | 0.0081 | 0.0092 | 0.0092 | 0.0085 | |
| 9/5/97 | 1501-1800 | 0.0033 | | 0.0104 | | 0.0077 | | |
| 9/5/97 | 1824-0257 | 0.0080 | 0.0080* | 0.0181* | 0.0127 | 0.0127* | | |
| 9/5/97 | 1824-0257 | -** | | -** | | | | Sampling tube contents transferred from broken tube |
| 9/6/97 | 0308-0550 | 0.0103 | 0.0103* | 0.0232 | 0.0232* | 0.0097 | 0.0097* | |
| 9/6/97 | 0308-0550 | - | | - | | - | | Sampling tube contents expelled from tube desorber |
| 9/6/97 | 0610-0855 | 0.0041 | 0.0212* | 0.0075 | 0.0335* | 0.0056 | 0.0067* | |
| 9/6/97 | 0610-0855 | 0.0212 | | 0.0335 | | 0.0067 | | |
| 9/6/97 | 0900-1158 | 0.0245 | 0.0142 | -** | -** | -** | 0.0018* | |
| 9/6/97 | 0900-1158 | 0.0039 | | -** | | | | |
| 9/6/97 | 1202-1458 | 0.0030 | 0.0035 | 0.0085 | 0.0092 | 0.0076 | 0.0081 | |
| 9/6/97 | 1205-1458 | 0.0039 | | 0.0099 | | 0.0087 | | |
| 9/6/97 | 1502-1800 | 0.0038 | 0.0038* | 0.0113 | 0.0113* | 0.0103 | 0.0103* | |
| 9/6/97 | 1502-1800 | 0.0481 | | 0.1317 | | 0.1055 | | Very low tricycene |
| BANNING OCT 3-4, 1997 | | | | | | | | |
| 10/3/97 | 0600-0858 | 0.0275 | 0.0275* | 0.0241 | 0.0241* | 0.0232 | 0.0232* | |
| 10/3/97 | 0600-0858 | 0.0469 | | -** | | 0.0277 | | |
| 10/3/97 | 0900-1158 | 0.0203 | 0.0214 | 0.0099 | 0.0099* | 0.0144 | 0.0134 | Altered monoterpene profile |
| 10/3/97 | 0900-1158 | 0.0226 | | -** | | 0.0124 | | |

Table C-1.3. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Banning (continued).

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|-----------|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|-----------------------------|
| 10/3/97 | 1200-1458 | 0.0159 | 0.0159 | ** | ** | 0.0032 | 0.0042 | |
| 10/3/97 | 1200-1458 | 0.0160 | | ** | | 0.0053 | | |
| 10/3/97 | 1500-1800 | 0.0066 | 0.0066* | 0.0157 | 0.0157* | 0.0085 | 0.0085* | |
| 10/3/97 | 1500-1800 | 0.0225 | | ** | | 0.0137 | | Altered monoterpane profile |
| 10/3-4/97 | 1807-0258 | 0.0247 | 0.0318* | 0.0122 | 0.0321* | 0.0243 | 0.0244* | Altered monoterpane profile |
| 10/3-4/97 | 1807-0258 | 0.0318 | | 0.0321 | | 0.0244 | | |
| 10/4/97 | 0300-0553 | 0.0590 | 0.0583 | 0.0785 | 0.0769 | 0.0356 | 0.0326 | |
| 10/4/97 | 0300-0553 | 0.0575 | | 0.0754 | | 0.0296 | | |
| 10/4/97 | 0601-0858 | 0.0464 | 0.0437 | 0.0586 | 0.0570 | 0.0194 | 0.0183 | |
| 10/4/97 | 0601-0858 | 0.0410 | | 0.0554 | | 0.0173 | | |
| 10/4/97 | 0900-1158 | 0.0098 | 0.0094 | 0.0234 | 0.0255 | 0.0141 | 0.0135 | |
| 10/4/97 | 0900-1158 | 0.0091 | | 0.0276 | | 0.0129 | | |
| 10/4/97 | 1200-1458 | 0.0052 | 0.0049 | 0.0168 | 0.0172 | 0.0111 | 0.0109 | |
| 10/4/97 | 1200-1458 | 0.0047 | | 0.0176 | | 0.0107 | | |
| 10/4/97 | 1500-1800 | 0.0052 | 0.0157 | 0.0124 | 0.0131 | 0.0077 | 0.0077 | |
| 10/4/97 | 1500-1800 | 0.0262 | | 0.0139 | | 0.0077 | | |

*Used single value.

**Not detected.

Table C-1.4. Detailed Data from Measurements of Limonene, 1,8-Cineole and Camphor in Ambient Air at Los Angeles, North Main Street.

| Date | Sampling | Limonene ppbv | Limonene av, ppbv | 1,8-Cineole ppbv | 1,8-Cineole av, ppbv | Camphor ppbv | Camphor av, ppbv | Reason for discarding data |
|---|-----------|------------------|----------------------|---------------------|-------------------------|-----------------|---------------------|----------------------------|
| LOS ANGELES NORTH MAIN ST. AUG 22-23, 1997 | | | | | | | | |
| 8/22/97 | 0558-0855 | 0.3211 | 0.3381 | 0.1014 | 0.1201 | 0.0463 | 0.0467 | |
| 8/22/97 | 0558-0855 | 0.3551 | | 0.1388 | | 0.0471 | | |
| 8/22/97 | 0900-1155 | 0.1208 | 0.1200 | 0.0558 | 0.0525 | 0.0433 | 0.0447 | |
| 8/22/97 | 0900-1155 | 0.1192 | | 0.0493 | | 0.0461 | | |
| 8/22/97 | 1202-1457 | 0.0189 | 0.0358 | 0.0173 | 0.0177 | 0.0091 | 0.0138 | |
| 8/22/97 | 1202-1457 | 0.0527 | | 0.0181 | | 0.0185 | | |
| 8/22/97 | 1500-1755 | 0.0244 | 0.0252 | 0.0059 | 0.0074 | 0.0050 | 0.0054 | |
| 8/22/97 | 1500-1755 | 0.0260 | | 0.0089 | | 0.0058 | | |
| 8/22-23/97 | 1806-0556 | 0.2304 | 0.2302 | 0.0360 | 0.0387 | 0.0188 | 0.0194 | |
| 8/22-23/97 | 1806-0556 | 0.2301 | | 0.0413 | | 0.0200 | | |
| 8/23/97 | 0605-0857 | 0.2299 | 0.2547 | 0.0511 | 0.0508 | 0.0243 | 0.0266 | |
| 8/23/97 | 0605-0857 | 0.2795 | | 0.0505 | | 0.0290 | | |
| 8/23/97 | 0900-1156 | 0.0312 | 0.0325 | 0.0262 | 0.0252 | 0.0157 | 0.0145 | |
| 8/23/97 | 0900-1156 | 0.0338 | | 0.0242 | | 0.0133 | | |
| 8/23/97 | 1202-1458 | 0.0072 | 0.0097 | 0.0077 | 0.0127 | 0.0058 | 0.0085 | |
| 8/23/97 | 1202-1458 | 0.0121 | | 0.0177 | | 0.0113 | | |
| 8/23/97 | 1500-1800 | 0.0057 | 0.0056 | -* | -* | 0.0034 | 0.0036 | |
| 8/23/97 | 1500-1800 | 0.0055 | | -* | | 0.0037 | | |

* Not detected.